

## A Comparative Analysis of Two High-Intensity Interval Training (HIIT) Programs on PGC-1 $\alpha$ , p53, and Citrate Synthase Protein Levels in Cardiomyocytes of Male Type 2 Diabetic Rats

Nadia Khayampour<sup>1</sup> , Maghsoud Peeri<sup>1</sup> , Mohammad Ali Azarbayjani<sup>1</sup> 

<sup>1</sup> Department of Exercise Physiology, Physical Education Faculty, Central Tehran Branch, Islamic Azad University, Tehran, Iran

### Article Info

#### Article type:

Research article

#### Article History:

Received: Feb. 23, 2022

Revised: Mar. 17, 2022

Accepted: Dec. 18, 2022

Published Online: Dec. 25, 2023

#### ✉ Correspondence to:

Maghsoud Peeri  
Department of Exercise  
Physiology, Central Tehran  
Branch, Islamic Azad  
University, Tehran, Iran

Email:  
[m.peeri@iauctb.ac.ir](mailto:m.peeri@iauctb.ac.ir)

### ABSTRACT

**Introduction:** This study investigates the impact of two high-intensity interval training (HIIT) programs on PGC-1 $\alpha$ , p53, and citrate synthase (CS) proteins within cardiomyocytes of male type 2 diabetic rats, aiming to discern potential molecular mechanisms influencing cardiac health.

**Material & Methods:** Twenty-four male Wistar rats were randomly assigned to control (NC), diabetic control (DC), diabetic with type 1 HIIT (HIIT-1), and diabetic with type 2 HIIT (HIIT-2) groups. Streptozotocin (STZ) induced type 2 diabetes, excluding the NC group. A four-week HIIT intervention, six sessions per week, preceded the analysis of heart tissue for PGC-1 $\alpha$ , p53, and CS protein levels. Statistical analysis employed GraphPad Prism software version 8 and one-way ANOVA ( $P < 0.05$ ).

**Results:** Both HIIT-1 ( $p=0.004$ ) and HIIT-2 ( $p=0.007$ ) groups exhibited significantly elevated cardiac PGC-1 $\alpha$  levels compared to DC. CS levels increased notably in HIIT-1 ( $p=0.001$ ) and HIIT-2 ( $p<0.001$ ), with HIIT-2 surpassing HIIT-1 significantly ( $p=0.010$ ). Concurrently, p53 levels significantly decreased in both HIIT-1 ( $p=0.005$ ) and HIIT-2 ( $p=0.001$ ) groups compared to DC.

**Conclusion:** Exercise training (HIIT) positively influences cardiac metabolism, evident in PGC-1 $\alpha$  and CS upregulation and p53 downregulation. While these findings provide valuable insights, further exploration is crucial for a comprehensive understanding of the underlying molecular mechanisms. This study advances our understanding of optimizing exercise interventions for improved cardiac health in type 2 diabetes.

**Keywords:** Exercise Therapy, Type 2 Diabetes Mellitus, Mitochondrial Diseases, PGC-1 $\alpha$ , Tumor Suppressor Protein p53, Citrate (si)-Synthase

### ➤ How to cite this paper

Khayampour N, Peeri M, Azarbayjani MA. A Comparative Analysis of Two High-Intensity Interval Training (HIIT) Programs on PGC-1 $\alpha$ , p53, and Citrate Synthase Protein Levels in Cardiomyocytes of Male Type 2 Diabetic Rats. J Bas Res Med Sci. 2023; 10(4):54-66.



© The Author(s)

Publisher: Ilam University of Medical Sciences

Journal of Basic Research in Medical Sciences: Volume 10, Issue 4, 2023

## Introduction

Type 2 diabetes mellitus (T2DM) is acknowledged as a heterogeneous condition arising from the interplay between non-modifiable genetic factors and modifiable environmental influences (1). Traditionally labeled as non-insulin dependent diabetes or adult-onset diabetes, T2DM is characterized by insulin resistance, which may progressively evolve into absolute resistance. However, in the past decade, reduced  $\beta$ -cell function has emerged as a pivotal issue in T2DM (2). Deviations in energy metabolism, including heightened fatty acid metabolism and diminished glucose metabolism, have been documented in individuals with diabetes, leading to a reduction in overall energy production (3). The accumulation of free fatty acids (FFA) in the cardiac tissue and the concomitant decrease in insulin-mediated glucose uptake among diabetic patients contribute to elevated cardiac oxygen consumption and mitochondrial dysfunction. These factors, in turn, precipitate cardiomyocyte death and result in ventricular dysfunction (4, 5).

Emerging evidence suggests that diabetes adversely affects mitochondrial function in the heart, thereby contributing to the development of diabetic cardiomyopathy (DCM) (6). Mitochondrial dysfunction and a reduction in mitochondrial biogenesis have been documented in diabetic patients, with researchers proposing that mitochondrial dysfunction may be implicated as a causative factor in insulin resistance (7). Among the key regulators of the mitochondrial biogenesis process and the expression of genes involved in oxidative phosphorylation is peroxisome proliferator-activated receptor gamma

coactivator 1 alpha (PGC-1 $\alpha$ ) (8). PGC-1 $\alpha$ , belonging to a small family of transcriptional coactivators, functions as a metabolic sensor that enables the body to respond to various stimuli such as exercise, fasting, and changes in metabolic substrate availability. Its dysregulation in the context of heart failure has garnered considerable attention (9). Existing evidence indicates that the induction of T2DM in rats through a high-fat diet combined with streptozotocin (STZ) injection is associated with a significant reduction in PGC-1 $\alpha$  levels in heart tissue. This reduction, in turn, leads to impaired cardiac mitochondrial function and biogenesis (10).

In spite of the crucial role played by PGC-1 $\alpha$  in the promotion of mitochondrial biogenesis, p53 assumes a significant function in the downregulation of PGC-1 $\alpha$  and the induction of cardiomyocyte necrosis (11, 12). Acknowledged as a prominent tumor suppressor, p53 emerges as a pivotal contributor to heart failure development (13), exerting regulatory control over diverse cellular functions. Elevated p53 expression fosters apoptosis and holds regulatory significance in cardiovascular health and disease (14). Within the nucleus, active p53 binds to the promoter region, thereby transactivating numerous target genes associated with cell cycle progression, apoptosis, and metabolism (12, 15). Investigations have delineated an augmented expression of p53 in type 2 diabetic rat cardiomyocytes compared to their healthy counterparts, and this heightened p53 expression correlates significantly with a reduction in PGC-1 $\alpha$  and citrate synthase (CS) expression (16). CS, a pivotal regulatory enzyme in the energy-generating metabolic pathway, catalyzes the condensation of

oxaloacetate and acetyl coenzyme A to generate citrate in the tricarboxylic acid cycle. It serves as a metabolic marker for assessing oxidative and respiratory capacity (17). Studies have reported diminished CS activity and protein levels of subunits from complexes I and III of the respiratory chain in obese insulin-resistant subjects compared to healthy subjects, underscoring the dysregulation of CS as a contributory factor to insulin resistance and the pathogenesis of diabetes (18).

A sedentary lifestyle, known as the cause of various diseases such as cancer and diabetes, underscores the importance of regular physical activity in promoting overall health and preventing these conditions (19-22). The positive effects of exercise training in preventing cardiac injury and enhancing mitochondrial biogenesis in advanced diabetic cardiomyopathy are linked to the activation of PGC-1 $\alpha$  and Akt (23). Despite the favorable impact of various forms of exercise training on diabetes management, a previous study indicated that High-Intensity Interval Training (HIIT) produced comparable, if not superior, improvements in body composition, physical fitness, and glycemic control when compared to continuous endurance training (22, 24). Consequently, HIIT is posited as a time-efficient therapeutic approach for individuals with Type 2 Diabetes (24-26). However, the underlying mechanisms of HIIT effectiveness remain substantially unknown. The current study is undertaken to compare the impact of two types of HIIT programs on the protein levels of PGC-1 $\alpha$ , p53, and citrate synthase in the

cardiomyocytes of male Type 2 diabetic rats.

## Materials and methods

### Animal

This semi experimental study, conducted on the 24 male Wistar rat (age ranging 5 to 6 weeks, 280 to 350 g weight) purchase from Razi Pasteur institutes. The rats were kept under a 12 h light-dark cycle (temperature:  $22 \pm 3$  C°) in tarbiat modares university animal house in similar environment and condition. The rats randomly assigned in four equal groups (6 rat in each group) including normal healthy control (NC), diabetic control (DC), type1 (HIIT-1) and type2 (HIIT-2) HIIT protocol. The animal in all groups had ad libitum access to chow and water throughout intervention period.

### Inducing type 2 diabetes (T2DM)

T2DM induced in the rats in all groups except NC group. For this purpose, following 12 hours night fasting the nicotinamide solution (110 mg/kg) were injected intraperitoneally. After 15 minutes, streptozotocin (STZ) freshly dissolved in (pH 4.5) citrate buffer injected intraperitoneally (60 mg/kg), dissolved citrate (0.05 mol) was gavage (12). The 72 hours after nicotinamide+STZ injection and in order to T2DM confirmation, the blood was collected from the tip of the tail vein for measurement of fasting blood glucose by glucometer, and blood glucose more than 200 mg/dl considered as T2DM. The rats body weight and fasting glucose levels in the different group reported in table1

**Table 1.** Body weight changes and glucose level in different group (mean $\pm$ SD).

Variable	NC	DC	HIIT-1	HIIT-2
Baseline weight (g)	318.3 $\pm$ 22.7	322.7 $\pm$ 11.3	328.7 $\pm$ 18.3	321.2 $\pm$ 18.0
Final weight (g)	376.2 $\pm$ 33.8	282.5 $\pm$ 39.3	268.7 $\pm$ 42.5	301.1 $\pm$ 55.3

Glucose/BG (mg/dl)	185.5 $\pm$ 12.7	567.3 $\pm$ 62.5	491.5 $\pm$ 33.4	427.5 $\pm$ 53.1
--------------------	------------------	------------------	------------------	------------------

NC: Healthy control, DC: Diabetic control, HIIT: High intensity interval training, BG: Blood Glucose.

### **High intensity interval training (HIIT) program**

Rats were familiarized with treadmills before exercise training began (10-15 min per day) with the intensity of 6 m/min for one week. Firstly, aerobic capacity of animals determined as previously reported procedure (27). The conducted exercise training program including two types of HIIT, both of them consist of three min warm up and three min cooldown with 40 % and 30% of rat maximum speed respectively. The type 1 HIIT protocol (HIIT-1) consist of 4 or 5 intensive intervals in the two first (1-2) and two final (3-4) weeks of training program respectively. High intensity intervals in

the first week, second week and 3-4 weeks conducted with 80%, 85% and 90% of determined rat maximum speed respectively. Each intensive interval duration was 2 min followed by 2 min low intensity interval with 40% of maximum speed. The type 2 HIIT protocol (HIIT-2) conducted with 5 intensive intervals in the first two weeks and 7 intervals in the two final weeks. Each intensive interval duration was 2 min followed by 2 min low intensity interval with 30% of maximum speed. The intensity of HIIT-2 protocol was similar to HIIT-1 protocol and covering distance of both HIIT protocol was similar. The both training program Properties have been reported in the table 2.

**Table 2.** Four weeks of HIIT program

Variable	Training	First week	Second week	Third week	Fourth week
Maximum speed when reaching VO <sub>2</sub> max (m/min)	HIIT-1	15	18	20	20
	HIIT-2	12	6	18	18
Training session time (min)	HIIT-1	6	8	10	12
	HIIT-2	6	8	10	12
Intensity (m/min)	HIIT-1	12	16	18	18
	HIIT-2	9	10	10	12

### **Blood sampling and tissue retrieval**

After a 48-hour fasting period following the conclusion of the final training session, animals were subjected to anesthesia using xylazine (10 mg/kg body weight, I.P.) and ketamine (90 mg/kg body weight, I.P.). Heparin-containing tubes were employed to collect blood samples from the hearts. After centrifugation at

15 °C and 3000 rpm for 15 minutes, serum was isolated and preserved at -80 °C for subsequent analysis. left ventricle were promptly harvested, flash-frozen in liquid nitrogen, and stored at -80 °C for future experimentation.

### **Blood glucose assessment**

For the determination of fasting blood Glucose (BG), the glucose oxidase

technique was employed, employing a quantitative glucose assay kit (Pars Azmoon, Karaj, Iran).

### ***Western blotting***

Western blot analysis was conducted on left ventricle cellular proteins, with 70-100 mg of tissue homogenized in RIPA buffer (pH 7.4, 1% Triton X-100, 50 mM Tris-HCl, 0.2% SDS, 0.2% sodium deoxycholate, 1 mM Na-EDTA, and 1 mM PMSF). The protein extracts, treated with PMSF and a protease inhibitor cocktail (Roche, Mannheim, Germany), were examined for changes in protein abundance. Total protein concentrations were determined using the Bradford assay. Subsequently, equal protein amounts from each sample were separated via SDS-PAGE, transferred to PVDF membranes, and blocked with 5% non-fat dry milk or bovine serum albumin in tris-buffered saline with 0.5% Tween-20 for two hours at room temperature. Primary antibodies targeting PGC-1 $\alpha$  (Cell signaling, Beverly, USA), p53 (SantaCruz, California, USA), CS (Cell signaling, Beverly, USA), and GAPDH (Cell signaling, Beverly, USA) were used for overnight incubation at 4 °C. Visualization of target protein bands occurred with an enhanced chemiluminescent substrate following

incubation with second HRP-conjugated antibodies (horseradish peroxidase). Band densities were analyzed using Image-J for densitometry (23).


### ***Statistical analysis***

The data obtained in the present study were subjected to analysis using GraphPad Prism software version 8. The normality of the data was assessed using the Shapiro-Wilk test, which indicated that the data followed a normal distribution. Between-group differences were assessed using a one-way ANOVA test, followed by the Tukey post hoc test for pairwise comparisons. Statistical significance was considered at the  $P < 0.05$  level.

### ***Results***

In the present study, all animals (normal and diabetic male Wistar rat) completed considered intervention (healthy or diabetic control, and HIIT program) and included in the final data analysis. One-way ANOVA test represented a significant between group difference for cardiac PGC-1 $\alpha$  ( $p < 0.001$ ), CS ( $p = 0.009$ ) and p53 ( $p < 0.001$ ) levels. Therefore, Tukey post hoc test was used for comparing different groups together.






**Figure 1.** PGC-1 $\alpha$  protein levels in cardiomyocyte. \* Significant difference with NC group. & Significant difference with DC group.

Present study findings indicated that cardiac PGC-1 $\alpha$  levels decreased significantly in the DC, HIIT-1 and HIIT-2 groups compared to NC group ( $p<0.05$ ). The levels of PGC-1 $\alpha$  in the HIIT-1 ( $p=0.004$ ) and HIIT-2 ( $p=0.007$ ) groups was significantly


higher compared to DC group. However, there was no significant difference between two trained groups (HIIT-1 and HIIT-2) for observed changes in the PGC-1 $\alpha$  levels ( $p=0.585$ ) (Figure 1).



**Figure 2.** Citrate synthase (CS) protein levels in cardiomyocyte. \* Significant difference with NC group. & Significant difference with DC group. # Significant difference with HIIT-1 group

The CS protein levels indicated a significant decrease in the DC, HIIT-1 and HIIT-2 groups compared to NC group ( $p < 0.05$ ). The CS protein levels in the HIIT-1 ( $p = 0.001$ ) and HIIT-2 ( $p < 0.001$ ) groups significantly increased compared to DC group. In addition, significant difference was

observed for CS protein between two trained groups (HIIT-1 and HIIT-2) and CS protein level in the HIIT-2 compared to HIIT-1 group was significantly higher ( $p = 0.010$ ), emphasized the further effect of HIIT-2 on CS protein (Figure 2).



**Figure 3.** p53 protein levels in cardiomyocyte. \* Significant difference with NC group. & Significant difference with DC group

Tukey post hoc test indicated a significant increase in cardiac p53 levels in the DC, HIIT-1 and HIIT-2 groups compared to NC group ( $p < 0.05$ ). On the other hand, significant decrease in p53 protein levels were observed in the HIIT-1 ( $p = 0.005$ ) and HIIT-2 ( $p = 0.001$ ) groups compared to DC group. However, there was no significant difference between HIIT-1 and HIIT-2 groups for observed changes in the cardiac p53 levels ( $p = 0.990$ ) (Figure 3).

## Discussion

The present study main findings were that four weeks HIIT in type 2 diabetic rat result in significant increase in PGC-1 $\alpha$  and CS proteins levels in the cardiomyocyte, and there was no significant difference between two exerted HIIT protocol for observed changes in PGC-1 $\alpha$  levels, but CS increase in HIIT-2 group was significant compared to HIIT-1 group. Yan et al (2013) reported that impairment of AMPK-PGC-1 $\alpha$  signaling results in mitochondrial biogenesis dysfunction, and this is a novel mechanism responsible for the

enhanced susceptibility of diabetic hearts to myocardial infarction (MI) and interventions that restore this signaling axis may have therapeutic potential for treating diabetic heart injury (28). In fact, pharmacological targets and drugs that modulate mitochondrial biogenesis, considered as a therapeutic strategy for the treatment of type 2 diabetes and insulin resistance (29).

Exercise training known as important stimulator for inducing mitochondrial biogenesis by activating different mechanism including PGC-1 $\alpha$  pathway (30), which present result supported this statement. Present findings regarding the PGC-1 $\alpha$  upregulation following exercise training confirmed in the previous study. wang et al (2020) reported that inducing diabetes with streptozotocin (STZ) injection in the male C57BL/6 mice is associated with decreasing cardiac PGC-1 $\alpha$  compared to non-diabetes mellitus sedentary control group (31). In contrast, these researchers reported that eight weeks endurance training on mouse treadmill result in significant increasing in cardiac PGC-1 $\alpha$  and AMPK protein expression and concluded that exercise training can ameliorates cardiac dysfunction by decreasing ROS production, enhances the mitochondrial oxidative capacity in the diabetic heart and exercise training shifts energy metabolism from fatty acid oxidation to glucose oxidation (31). In addition, SIRT1/PGC-1 $\alpha$ /PI3K/Akt pathway downregulation after MI, restores by post-MI 4-week treadmill exercise training, and similar to MI group, activation of SIRT1/PGC-1 $\alpha$ /PI3K/Akt signaling pathway were observed in healthy adult male rats (32).

In addition, some researchers attributed the protective effects of

exercise training (three weeks swimming) against acute rodent model myocardial infarction to improving myocardial energy metabolism and mitochondrial biogenesis by an activation of PGC-1 $\alpha$  in heart tissue, which cardiac PGC-1 $\alpha$  upregulation by swimming training was associated with significant increase in expression of other genes involved in mitochondrial biogenesis such as TFAM, NRF-1 and NRF-2, and simultaneous decrease in apoptotic genes including caspase 3 and BAX were observed (33). Unfortunately, in the present study the changes in the above-mentioned genes don't investigate. However, we observed that cardiac p53 protein as an important apoptosis stimulator decreased in diabetic rat after four weeks HIIT. Studies have reported that p53 and associated apoptosis play important roles in the pathogenesis of ischemia/reperfusion caused acute cardiac diseases (34). p53 promotes cardiac dysfunction in diabetes via excessive mitochondrial respiration-mediated oxidative stress and lipid accumulation (35). On the other hand, inhibition of p53 prevents diabetic cardiomyopathy by preventing early-stage apoptosis and cell senescence (36), and according to our findings high intensity interval training is effective strategy for p53 downregulation. Al-Jarrah et al (2012) supported our findings and suggested that levels of p53 increased in the heart of diabetic rats, and 4-week treadmill training program significantly decreased the expression level of p53 in in the cardiac muscle (37). Moreover, decreased in p53 protein levels have been observed after 8 weeks endurance training on treadmill in other tissue including skeletal muscle of type 2 diabetic Goto-Kakizaki rats, which result in attenuates oxidative stress and researchers concluded that p53



downregulation is an important mechanism for exercise-induced increase in mitochondrial content and function (38). Since, elevated p53 levels correlate with cardiomyocyte apoptosis and impairment of mitochondrial function (39), its inhibition plays an important role for trigger the exercise training induced adaption in the heart tissue. Although, the changes in oxidative stress markers and mitochondrial content and function in our research is not studied, but despite observed decrease in the p53 levels and simultaneous increase in CS and PGC-1 $\alpha$  levels as mitochondrial biogenesis stimulators, it can be concluded that mitochondrial biogenesis probably increased in the present study.

In corroborate with our findings regarding CS upregulation in cardiac tissue by exercise training, Siu et al (2003) reported that eight weeks aerobic training in Sprague-Dawley rats cause to significant enhancement of expression and enzyme activity of citrate synthase in cardiac and skeletal (soleus) muscles (40). Moreover, increase in CS activity in untrained and trained human skeletal muscle have been reported, although the mechanism of this increase is unknown (41). Some researchers observed enhancing CS activity following different type of exercise training including HIIT and showed a positive and significant linear relationship between CS activity and VO<sub>2</sub>max (42). Collectively, present study findings suggested that HIIT protocol may result in improving cardiac metabolism through CS and PGC-1 $\alpha$  upregulation as a stimulator of mitochondrial biogenesis and decreasing p53 level which play important role in cardiomyocyte apoptosis and impairment of mitochondrial function.

## Conclusion

In conclusion, this study sheds light on the positive impact of four weeks of HIIT on cardiac tissue in type 2 diabetic rats. The observed increases in PGC-1 $\alpha$  and CS protein levels, coupled with a decrease in p53, suggest potential mechanisms contributing to improved mitochondrial function and biogenesis. These findings align with the broader literature on exercise-induced adaptations and hold promise for addressing cardiac dysfunction in diabetes. However, several limitations, including the focused protein-level analysis, the absence of exploration into broader molecular markers, and the reliance on a rat model, should be acknowledged. While our results provide valuable insights, further research is warranted to delve into the long-term effects, individual-specific responses, and broader molecular changes associated with HIIT. Understanding these aspects will not only refine our comprehension of the mechanisms at play but also enhance the translatability of these findings to human populations. In essence, the current study sets the stage for future investigations to unravel the full spectrum of benefits and considerations associated with incorporating HIIT as a potential therapeutic strategy for mitigating cardiac complications in type 2 diabetes.

## Acknowledgements

This research was extracted from the findings of a Ph.D. thesis in exercise physiology. The authors would like to express their gratitude to all individuals who contributed to the implementation of this study.

## Financial support

No financial support was received for this research.

**Conflict of interest**

The authors declare that no conflict of interest exists.

**Authors' contributions**

N Kh. conducted the study protocol, M. P. designed and confirmed the training protocol, and M A A. analyzed the research data. The article draft was prepared collaboratively by N Kh. and M A A., and the final manuscript was edited by M P. All authors contributed equally to data collection.

## References

1. Doria A, Patti M-E, Kahn CR. The emerging genetic architecture of type 2 diabetes. *Cell metabolism*. 2008;8(3):186-200. doi: 10.1016/j.cmet.2008.08.006.
2. Artasensi A, Pedretti A, Vistoli G, Fumagalli L. Type 2 diabetes mellitus: a review of multi-target drugs. *Molecules*. 2020;25(8):1987. doi: 10.3390/molecules25081987.
3. Miki T, Yuda S, Kouzu H, Miura T. Diabetic cardiomyopathy: pathophysiology and clinical features. *Heart failure reviews*. 2013;18(2):149-66. doi: 10.1007/s10741-012-9313-3.
4. Rijzewijk LJ, van der Meer RW, Lamb HJ, de Jong HW, Lubberink M, Romijn JA, et al. Altered myocardial substrate metabolism and decreased diastolic function in nonischemic human diabetic cardiomyopathy: studies with cardiac positron emission tomography and magnetic resonance imaging. *Journal of the American College of Cardiology*. 2009;54(16):1524-32. doi: 10.1016/j.jacc.2009.04.074.
5. Khakdan S, Delfan M, Heydarpour Meymeh M, Kazerouni F, Ghaedi H, Shanaki M, et al. High-intensity interval training (HIIT) effectively enhances heart function via miR-195 dependent cardiomyopathy reduction in high-fat high-fructose diet-induced diabetic rats. *Archives of physiology and biochemistry*. 2020;126(3):250-7. doi: 10.1080/13813455.2018.1511599
6. Bombicino SS, Iglesias DE, Mikusic IAR, D'Annunzio V, Gelpi RJ, Boveris A, et al. Diabetes impairs heart mitochondrial function without changes in resting cardiac performance. *The international journal of biochemistry & cell biology*. 2016; 81:335-45. doi: 10.1016/j.biocel.2016.09.018.
7. Wang CH, Wang CC, Wei YH. Mitochondrial dysfunction in insulin insensitivity: implication of mitochondrial role in type 2 diabetes. *Annals of the New York Academy of Sciences*. 2010;1201(1):157-65. doi: 10.1111/j.1749-6632.2010.05625.x.
8. Puigserver P, Spiegelman BM. Peroxisome proliferator-activated receptor- $\gamma$  coactivator 1 $\alpha$  (PGC-1 $\alpha$ ): transcriptional coactivator and metabolic regulator. *Endocrine reviews*. 2003;24(1):78-90. doi: 10.1210/er.2002-0012. doi: 10.3389/fcvm.2020.00002.
9. Oka S-i, Sabry AD, Cawley KM, Warren JS. Multiple levels of PGC-1 $\alpha$  dysregulation in heart failure. *Frontiers in Cardiovascular Medicine*. 2020;2.
10. Fang W-j, Wang C-j, He Y, Zhou Y-l, Peng X-d, Liu S-k. Resveratrol alleviates diabetic cardiomyopathy in rats by improving mitochondrial function through PGC-1 $\alpha$  deacetylation. *Acta Pharmacologica Sinica*. 2018;39(1):59-73. doi: 10.1038/aps.2017.50.
11. Villeneuve C, Guilbeau-Frugier C, Sicard P, Lairez O, Ordener C, Duparc T, et al. p53-PGC-1 $\alpha$  pathway mediates oxidative mitochondrial damage and cardiomyocyte necrosis induced by monoamine oxidase-A upregulation: role in chronic left ventricular dysfunction in mice. *Antioxidants & redox signaling*. 2013;18(1):5-18. doi: 10.1089/ars.2011.4373.
12. Delfan M, Vahed A, Bishop DJ, Amadeh Juybari R, Laher I, Saeidi A, et al. Effects of two workload-matched high intensity interval training protocols on regulatory factors associated with mitochondrial biogenesis in the soleus muscle of diabetic rats. *Frontiers in Physiology*. 2022; 13:927969. doi: 10.3389/fphys.2022.927969
13. Fujita T, Ishikawa Y. Apoptosis in Heart Failure-The Role of the  $\beta$ -Adrenergic Receptor-Mediated Signaling Pathway and p53-Mediated Signaling Pathway in the Apoptosis of Cardiomyocytes-. *Circulation Journal*. 2011;75(8):1811-8. doi: 10.1253/circj. cj-11-0025.
14. Men H, Cai H, Cheng Q, Zhou W, Wang X, Huang S, et al. The regulatory roles of p53 in cardiovascular health and disease. *Cellular and Molecular Life Sciences*. 2021;78(5):2001-18. doi: 10.1007/s00018-020-03694-6.
15. Vousden KH, Lane DP. p53 in health and disease. *Nature reviews Molecular cell biology*. 2007;8(4):275-83. doi: 10.1038/nrm2147.
16. Khayampour N, Peeri M, Azarbayjani MA, Delfan M. Effects of High Intensity Interval Training on the Gene Expression of PGC1-A, CS and P-53 in the Cardiomyocyte of Male Obese Rats in Type 2 Diabetes. *Journal of Shahid Sadoughi University of Medical Sciences*. 2020. doi: 10.18502/ssu.v28i11.5222. doi: 10.18502/ssu.v28i11.5222.
17. Spina RJ, Chi M, Hopkins MG, Nemeth P, Lowry O, Holloszy J. Mitochondrial enzymes increase in muscle in response to 7-10 days of cycle exercise. *Journal of applied physiology*. 1996;80(6):2250-4. doi: 10.1152/jappl.1996.80.6.2250.

18. Heilbronn LK, Gan SK, Turner N, Campbell LV, Chisholm DJ. Markers of mitochondrial biogenesis and metabolism are lower in overweight and obese insulin-resistant subjects. *The Journal of Clinical Endocrinology & Metabolism*. 2007;92(4):1467-73. doi: 10.1210/jc.2006-2210.
19. Teixeira-Lemos E, Nunes S, Teixeira F, Reis F. Regular physical exercise training assists in preventing type 2 diabetes development: focus on its antioxidant and anti-inflammatory properties. *Cardiovascular diabetology*. 2011;10(1):1-15. doi: 10.1186/1475-2840-10-12.
20. Delfan M, Delphan M, Kordi MR, Ravasi AA, Safa M, Gorgani-Firuzjaee S, et al. High intensity interval training improves diabetic cardiomyopathy via miR-1 dependent suppression of cardiomyocyte apoptosis in diabetic rats. *Journal of Diabetes & Metabolic Disorders*. 2020; 19:145-52. doi: 10.1007/s40200-019-00485-0
21. Mirakhori Z, Kordi MR, Alizadeh S, Anoosheh L, Amani Shalamzari S, Amini A, et al. The effect of aerobic training on plasma estradiol and mir-206 and  $\text{er}\alpha$  expression in mice with breast cancer. *Iranian Journal of Breast Diseases*. 2015;7(4):23-32. doi: 10.1001.1.17359406.1393.7.4.3.7
22. Akbari N, Peeri M, Azarbayjani MA, Delfan M. Comparison of the effect of 8 weeks of continuous and high intensity interval training on the gene expression of TIMP-2 and MMP-2 in male diabetic rats. *Razi Journal of Medical Sciences*. 2019;26(10):107-16. <http://rjms.iums.ac.ir/article-1-5702-en.html>
23. Wang H, Bei Y, Lu Y, Sun W, Liu Q, Wang Y, et al. Exercise prevents cardiac injury and improves mitochondrial biogenesis in advanced diabetic cardiomyopathy with PGC-1 $\alpha$  and Akt activation. *Cellular physiology and biochemistry*. 2015;35(6):2159-68. doi: 10.1159/000374021.
24. Ghafari S, Nazarali P, Razavi A, Delfan M. Effect of continuous aerobic training versus high intensity interval training on Resistin and insulin resistance in type 2 diabetic rats. *Journal of Shahid Sadoughi University of Medical Sciences*. 2019.
25. Winding KM, Munch GW, Iepsen UW, Van Hall G, Pedersen BK, Mortensen SP. The effect on glycaemic control of low-volume high-intensity interval training versus endurance training in individuals with type 2 diabetes. *Diabetes, Obesity and Metabolism*. 2018;20(5):1131-9. doi: 10.1111/dom.13198.
26. Rezaee N, Rahmani-Nia F, Delfan M, Ghahremani R. Exercise training and probiotic supplementation effects on skeletal muscle apoptosis prevention in type-I diabetic rats. *Life Sciences*. 2021; 285:119973. doi: 10.1016/j.lfs.2021.119973
27. PITHON-CURI TNC. Aprogram Of Moderate Physical Training For Wistar Rats Based On Maximal Oxygen Consumption. *J Strength Cond Res*. 2007; 21(3):751-6. doi: 10.1519/R-20155.1.
28. Yan W, Zhang H, Liu P, Wang H, Liu J, Gao C, et al. Impaired mitochondrial biogenesis due to dysfunctional adiponectin-AMPK-PGC-1 $\alpha$  signaling contributing to increased vulnerability in diabetic heart. *Basic Res Cardiol*. 2013; 108(3):1-15. doi: 10.1007/s00395-013-0329-1.
29. Zamora M, Pardo R, Villena JA. Pharmacological induction of mitochondrial biogenesis as a therapeutic strategy for the treatment of type 2 diabetes. *Biochem Pharmacol*. 2015; 98(1):16-28. doi: 10.1016/j.bcp.2015.06.032.
30. Hood DA. Mechanisms of exercise-induced mitochondrial biogenesis in skeletal muscle. *Applied Physiology, Nutrition, and Metabolism*. 2009;34(3):465-72.
31. Wang SY, Zhu S, Wu J, Zhang M, Xu Y, Xu W, et al. Exercise enhances cardiac function by improving mitochondrial dysfunction and maintaining energy homeostasis in the development of diabetic cardiomyopathy. *J Mol Med*. 2020; 98(2):245-61. doi: 10.1007/s00109-019-01861-2.
32. Jia D, Hou L, Lv Y, Xi L, Tian Z. Postinfarction exercise training alleviates cardiac dysfunction and adverse remodeling via mitochondrial biogenesis and SIRT1/PGC-1 $\alpha$ /PI3K/Akt signaling. *J Cell Physiol*. 2019; 234(12):23705-18. doi: 10.1002/jcp.28939.
33. Tao L, Bei Y, Lin S, Zhang H, Zhou Y, Jiang J, et al. Exercise training protects against acute myocardial infarction via improving myocardial energy metabolism and mitochondrial biogenesis. *Cell Physiol Biochem*. 2015; 37(1):162-75. doi: 10.1159/000430342.
34. Zhang Y, Köhler K, Xu J, Lu D, Braun T, Schlitt A, et al. Inhibition of p53 after acute myocardial infarction: reduction of apoptosis is counteracted by disturbed scar formation and cardiac rupture. *J Mol*

- Cell Cardiol. 2011; 50(3):471-8. doi: 10.1016/j.yjmcc.2010.11.006.
35. Nakamura H, Matoba S, Iwai-Kanai E, Kimata M, Hoshino A, Nakaoka M, et al. p53 promotes cardiac dysfunction in diabetic mellitus caused by excessive mitochondrial respiration-mediated reactive oxygen species generation and lipid accumulation. *Circ Heart Fail.* 2012; 5(1):106-15. doi: 10.1161/CIRCHEARTFAILURE.111.961565.
  36. Gu J, Wang S, Guo H, Tan Y, Liang Y, Feng A, et al. Inhibition of p53 prevents diabetic cardiomyopathy by preventing early-stage apoptosis and cell senescence, reduced glycolysis, and impaired angiogenesis. *Cell Death Dis.* 2018; 9(2):1-17. doi: 10.1038/s41419-017-0093-5.
  37. Al-Jarrah M, Ahmad MB, Maayah M, Al-Khatib A. Effect of exercise training on the expression of p53 and iNOS in the cardiac muscle of type I diabetic rats. *J Clin Endocrinol Metab.* 2012; 2(4-5):176-80. doi: 10.4021/jem123e.
  38. Qi Z, He J, Zhang Y, Shao Y, Ding S. Exercise training attenuates oxidative stress and decreases p53 protein content in skeletal muscle of type 2 diabetic Goto-Kakizaki rats. *Free Radic Biol Med.* 2011; 50(7):794-800. doi: 10.1016/j.freeradbiomed.2010.12.022.
  39. Mak TW, Hauck L, Grothe D, Billia F. p53 regulates the cardiac transcriptome. *Proc Natl Acad Sci.* 2017; 114(9):2331-6. doi: 10.1073/pnas.1621436114.
  40. Siu PM, Donley DA, Bryner RW, Alway SE. Citrate synthase expression and enzyme activity after endurance training in cardiac and skeletal muscles. *J Appl Physiol.* 2003;94(2):555-60. doi: 10.1152/japplphysiol.00821.2002.
  41. Leek BT, Mudaliar SR, Henry R, Mathieu-Costello O, Richardson RS. Effect of acute exercise on citrate synthase activity in untrained and trained human skeletal muscle. *Am J Physiol Regul Integr Comp Physiol.* 2001; 280(2): R441-R7. doi: 10.1152/ajpregu.2001.280.2. R441.
  42. Vigelsø A, Andersen NB, Dela F. The relationship between skeletal muscle mitochondrial citrate synthase activity and whole-body oxygen uptake adaptations in response to exercise training. *Int J Physiol Pathophysiol Pharmacol.* 2014; 6(2):84-101. PMID: 25057335.