

Risk factors affecting abortion among pregnant women– A case-control study

Shima Hashemi¹ , Ashraf Direkvand Moghadam² , Kourosh Sayehmiri³ , Mirhadi Mousavi⁴ , Hamzeh Ahmadi⁵ , Monireh Azizi⁶ , Khairollah Asadollahi¹  

¹ Department of Epidemiology, Faculty of Health, Ilam University of Medical Sciences, Ilam, Iran

² Department of Midwifery, Faculty of Nursery and Midwifery, Ilam University of Medical Sciences, Ilam, Iran

³ Department of Biostatistics, Faculty of Health, Ilam University of Medical Sciences, Ilam, Iran

⁴ Department of Pediatrics, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

⁵ Department of Epidemiology and Biostatistics, Faculty of Public Health, Isfahan University of Medical Sciences, Isfahan, Iran

⁶ Department of Anatomy, Faculty of Medicine, Ilam University of Medical Sciences, Ilam, Iran

Article Info

Article type:

Research article

Article History:

Received: Dec. 14, 2023

Accepted: Mar. 25, 2024

Published Online: Mar. 06, 2024

✉ Correspondence to:

Khairollah Asadollahi
Department of Epidemiology,
Faculty of Health, Ilam
University of Medical Sciences,
Ilam, Iran

Email:

masoud_1241@yahoo.co.uk

ABSTRACT

Introduction: Abortion, a common pregnancy complication, is influenced by multifaceted reproductive, economic, social, and cultural factors. This study investigates abortion determinants among pregnant women in Khorramabad, Iran, from September 2020 to March 2021.

Material & Methods: This case-control study involved 416 pregnant women (208 cases, 208 controls) selected from Khorramabad health centers through census and random sampling during the COVID-19 pandemic. Data on demographics and pregnancy history were collected using a checklist and analyzed with STATA16.

Results: Among the 416 pregnant women, the mean age was 31.00 ± 5.53 years in the control group and 32.27 ± 6.85 years in the case group. Furthermore, 74 (35.58%) mothers in the case group had a diploma, 192 (92.31%) were housewives, and 106 (50.96%) had A+ blood group. The majority of mothers (46.63% in case, 39.42% in control) had an overweight range of pre-pregnancy BMI (25-29.9) (kg/m²). The mean height was 162.97 ± 5.30 in the control group and 160.86 ± 5.39 in the case group. Multiple regression logistic analysis confirmed significant associations between abortion and mother's age (OR=4.67, 95%CI=1.11-19.62), previous cesarean delivery (OR=4.33, 95%CI=1.94-9.66), not taking folic acid pills during pregnancy (OR=14.92, 95%CI=6.25-35.61), low income (OR=7.00, 95%CI=2.02-24.35), not smoking (OR=0.28, 95%CI=0.08-0.94), and absence of pre-diabetes (OR=0.13, 95%CI=0.03-0.47) ($p < 0.05$). No significant relationships were found between abortion and other studied risk factors.

Conclusion: Older mothers with lower income levels are more likely to experience abortion. Additionally, factors such as pre-existing pre-diabetes, previous cesarean deliveries, smoking, lack of folic acid supplementation, and maternal infections are identified as risk factors for abortion. Providing education, healthcare, and continuous support from healthcare workers can promote safe pregnancies, reduce the risk of abortion, and enhance maternal and child health outcomes.

Keywords: Abortion, Pregnancy, Mothers, Child

➤ How to cite this paper

Hashemi Sh, Direkvand Moghadam A, Sayehmiri K, Mousavi M, Ahmadi H, Azizi M, Asadollahi Kh. Risk factors affecting abortion among pregnant women– A case-control study. J Bas Res Med Sci. 2024; 11(2):1-13.



© The Author(s)

Publisher: Ilam University of Medical Sciences

Introduction

Abortion is defined as fetal loss before 20 gestational weeks or ≤ 500 gram weight (1,2). Abortion is a significant social and health issue worldwide and one of the most complications in the pregnancy which affects women's life (3). In the last decade, the prevalence of spontaneous abortions has increased (4). Spontaneous abortion is defined as fetal loss before twenty gestational weeks which occurs in 10-20% of pregnancies. Many different genetics and physiological factors may cause spontaneous abortion (4,5). According to estimation, around 73 million induced abortions take place worldwide each year. Six out of 10 (61%) of all unintended pregnancies, and 3 out of 10 (29%) of all pregnancies, end in induced abortion (6).

During 2010 to 2014, over than 55.7 million abortions took place each year in the world. About 25.1 million abortions were unsafe and 97% of them occurred in developing countries (7). Annually, 121 million unwanted pregnancies have occurred in the world, which 61% of them resulted in abortion (8). Accurate statistics of abortions in Iran are not available, and according to researches, the number of illegal abortions is about 290,000 annually, and the highest age range for abortions is between 15 and 24 years old (9). Studies showed that there are some factors affecting abortion including: uterine abnormalities, immune system disorders, hormonal disorders, anatomical defects and endometriosis, mother's age, environmental factors, infections, maternal diseases such as genetic disorders and chromosomal abnormalities, endocrine disorders such as hypothyroidism and diabetes, medication use, smoking, alcohol, caffeine, exposure to mobile phone radiation, use of contraceptive drugs, trauma, blood factors and other unknown factors (10-12).

The harmful consequence of abortion on mother's reproductive capability is evident. Abortion leads to a high burden of complications including: bleeding, uterine infection, toxic shock syndrome, sepsis, acute kidney failure, parametrit, peritonitis, and

even maternal deaths (13). Abortion is a distressing experience that affects mothers in a variety of ways by influencing on emotional status and may lead to physical disorders, depression, anxiety, sleep disorders and general health complications and reduces the individual's social functioning (14). The consequences of unsafe abortion do not only affect the individual, but also affect the entire health care system because many resources must be spent on treating the complications caused by unsafe abortions (15).

Maternal mortality due to complications of pregnancy and childbirth is one of the most important indicators showing the level of development of countries and consists of a set of different factors, while preventive and appropriate interventions can prevent the occurrence of similar deaths in the future. Abortion is one of the pervasive issues in human societies, which not only requires comprehensive medical and health investigations, but is also the result of countless causes that can be investigated in different jurisprudential, legal and social fields. This study aimed to determine risk factors affecting abortion among pregnant women in a big city located in the west of Iran.

Materials and methods

In this case-control study, 416 pregnant women (208 cases, 208 controls) referred to Khorramabad health centers under the supervision of the Ministry of Health and Medical Education (MOHME) were selected by census sampling as the case group and random sampling as the control group, from September 2020 to March 2021. Written informed consent and verbal agreement was taken from all participants. All experimental protocols were approved by Ilam University of Medical Sciences.

There are 16 health centers in Khorramabad city. Women who had lost their pregnancy under 20 weeks of gestational age were defined as the case group, and women with ongoing pregnancies from 20 to 42 weeks of gestational age were defined as the control group. The sample size was calculated

based on the number of abortions reported during September 2020 to March 2021 among health centers using the formula below, with an equal number among non-aborted women totaling 416 participants. Women with a history of pregnancy were included in this study. The inclusion criteria for women were being eager to participate and complete the research questions. There was no limitation of age range. The authors excluded those who had no history of marriage and pregnancy, as well as those who had immigrated to other cities and had no printed or electronic health center documents.

$$n = \left[\frac{z_{1-\alpha/2} \sqrt{p_1(1-p_1)} + z_{1-\beta} \sqrt{p_1(1-p_1) + p_2(1-p_2)}}{(p_1 - p_2)} \right]^2 = 205$$

n (case): 205, n (control): 205, n (total): 410, P1: exposure in cases, P2: exposure in controls, Z1- $\alpha/2$ = 1.96 and Z1- β = 1.28, CI: 95%, Power: 80%, Population proportion: 2, OR: 2.16

The data were analyzed using STATA 16. P values ≤ 0.05 were considered statistically significant. Continuous variables were presented as mean \pm standard deviation, while qualitative variables were expressed as frequency (percent). To investigate the risk factors of abortion, a univariate logistic regression analysis was initially conducted. Variables with $p \leq 0.2$ in the univariate analysis were then included in the multiple regression model. Subsequently, variables with $p \leq 0.05$ were retained, while others were excluded using a stepwise backward method. The effects on risk were estimated by odds ratios with 95% confidence intervals, and p-values ≤ 0.05 were deemed statistically significant. Variables with a P value < 0.2 were further tested using multivariate analysis, employing backward-elimination multiple logistic regression. This process aimed to identify the most statistically significant variables as risk factors.

Data were collected using a checklist from the Iranian electronic health record information, known as the "SIB" Integrated Health System. The checklist included questions about demographic information, the participant's pregnancy history, the participant's medical history, and socio-economic information. Women with incomplete record information were invited to complete the checklist in person at the health centers where they were enrolled. Written informed consent was obtained from all participants. The checklist was anonymous, and the interviewers were unaware of the name or record number of participants. This study was extracted from an MSc Epidemiology thesis and was approved by Ilam University of Medical Sciences and the Ethics Committee with the approval number IR.MEDILAM.REC.1400.193. All methods were conducted in accordance with relevant guidelines, regulations, or the Declaration of Helsinki.

Results

Based on the results, among the 416 analyzed participants, the mean \pm standard deviation age was 31.00 ± 5.53 years in the control group and 32.27 ± 6.85 years in the case group. Additionally, the majority of mothers (74, 35.58%) had a diploma, 192 (92.31%) were housewives, and 106 (50.96%) had A+ blood group in the case group. It was also found that the majority of mothers (97, 46.63%) in the case group and 82 (39.42%) in the control group had an overweight range of pre-pregnancy BMI (25-29.9 kg/m²), respectively. From the total number of women in the control group, 189 (90.87%) had been taking folic acid pills one month before pregnancy. There was a significant association between not using folic acid and abortion ($p < 0.0001$). Demographic information is presented in Table 1.

Table 1. Comparison of Demographic Information in Mothers with and without Abortion.

| Variables | Category | Case group N (%) | Control group N (%) | P-value |
|--|---------------------------|---------------------|------------------------|---------|
| Mother educational level | Elementary to high school | 55 (26.44) | 51 (24.52) | 0.328 |
| | Diploma | 74 (35.58) | 65 (31.25) | |
| | Associated degree | 22 (10.58) | 16 (7.69) | |
| | BSc | 47 (22.60) | 65 (31.25) | |
| | MSc & PhD | 10 (4.80) | 11 (5.29) | |
| Mother job status | Housewife | 192 (92.31) | 185 (88.94) | 0.464 |
| | Employee | 13 (6.25) | 19 (9.14) | |
| | Self-employee | 3 (1.44) | 4 (1.92) | |
| Mother blood group | A+ | 106 (50.96) | 65 (31.25) | <0.0001 |
| | A- | 12 (5.77) | 1 (0.48) | |
| | B+ | 30 (14.42) | 23 (11.06) | |
| | B- | 1 (0.48) | 4 (1.92) | |
| | AB+ | 16 (7.69) | 10 (4.81) | |
| | AB- | 0 | 1 (0.48) | |
| | O+ | 40 (19.23) | 96 (46.15) | |
| O- | 3 (1.45) | 8 (3.85) | | |
| pre-pregnancy Body Mass Index (BMI) ($\frac{\text{kg}}{\text{m}^2}$) | ≤ 18.4 | 1(0.48) | 4(1.92) | 0.012 |
| | 18.5-24.9 | 51 (24.52) | 80 (38.64) | |
| | 25-29.9 | 97 (46.63) | 82 (39.42) | |
| | 30-34.9 | 48 (23.08) | 33 (15.86) | |
| | ≥ 35 | 11 (5.29) | 9 (4.34) | |
| Husband educational level | Elementary to high school | 54 (25.96) | 40 (19.23) | 0.131 |
| | Diploma | 93 (44.72) | 83 (39.90) | |
| | Associated degree | 17 (8.17) | 20 (9.62) | |
| | BSc | 31 (14.90) | 48 (23.08) | |
| | MSc & PhD | 13 (6.25) | 17 (8.17) | |
| Husband job status | Unemployed | 17 (8.17) | 10 (4.81) | 0.309 |
| | Employee | 56 (26.92) | 64 (30.77) | |
| | Self-employee | 135 (64.91) | 134 (64.42) | |
| Husband blood group | A+ | 41 (19.71) | 52 (25.0) | 0.029 |
| | A- | 37 (17.79) | 33 (15.87) | |
| | B+ | 25 (12.02) | 15 (7.21) | |
| | B- | 10 (4.81) | 2 (0.96) | |
| | AB+ | 26 (12.50) | 39 (18.75) | |
| | AB- | 3 (1.44) | 0 | |
| | O+ | 63 (30.29) | 62 (29.81) | |
| O- | 3 (1.44) | 5 (2.40) | | |
| Pregnancy type | Wanted | 165 (79.33) | 169 (81.25) | 0.622 |
| | Unwanted | 43 (20.67) | 39 (18.75) | |
| Last delivery type | Vaginal | 89(59.33) | 110(84.62) | <0.0001 |
| | Cesarean | 61(40.67) | 20(15.38) | |
| Contraceptive methods | Natural family planning | 94 (45.41) | 57 (27.40) | 0.002 |

| | | | | |
|--|-------------------------|-------------|-------------|---------|
| | Condom | 70 (33.81) | 107 (51.44) | |
| | Combined pills | 25 (12.08) | 27 (12.98) | |
| | Contraceptive injection | 9 (4.35) | 7 (3.37) | |
| | IUD | 9 (4.35) | 10 (4.81) | |
| Caffeine consumption | Yes | 159 (76.44) | 157 (75.48) | 0.819 |
| | No | 49 (23.56) | 51 (24.52) | |
| Cup of coffee (per day) | 0 | 41 (19.71) | 51 (24.52) | 0.009 |
| | 1 | 32 (15.38) | 56 (26.92) | |
| | 2 | 46 (22.12) | 36 (17.30) | |
| | 3 | 36 (17.31) | 35 (16.83) | |
| | 4 | 9 (4.33) | 9 (4.33) | |
| | 5 | 15 (7.21) | 8 (3.85) | |
| | 6 \geq | 29 (13.94) | 13 (6.25) | |
| Smoking | Yes | 47 (22.60) | 14 (6.73) | <0.0001 |
| | No | 161 (77.40) | 194 (93.27) | |
| Monthly income | \leq 150 \$ | 42 (20.19) | 18 (8.65) | <0.0001 |
| | 150-250 \$ | 112 (53.85) | 105 (50.49) | |
| | \geq 250 \$ | 54 (25.96) | 85 (40.86) | |
| Maternal Disorder | Yes | 17(8.17) | 4(1.92) | 0.004 |
| | No | 191(91.83) | 204(98.08) | |
| Maternal Disorder Type | Autoimmune disease | 3(15.79) | 3(75.0) | 0.047 |
| | Metabolic | 3(15.79) | 0 | |
| | Infections | 13(68.42) | 1(25.0) | |
| Acid folic (3 months before pregnancy) | Yes | 59(28.37) | 114(54.81) | <0.0001 |
| | No | 149(71.63) | 94(45.19) | |
| Supplementary (3months before pregnancy) | Yes | 22(10.58) | 120(57.69) | <0.0001 |
| | No | 186(89.42) | 88(42.31) | |
| Acid folic (1 months before pregnancy) | Yes | 96(46.15) | 189(90.87) | <0.0001 |
| | No | 112(53.85) | 19(9.13) | |
| Drug Consumption | Yes | 68(32.69) | 73(35.10) | 0.605 |
| | No | 140(67.31) | 135(64.90) | |
| Levothyroxine Use | Yes | 24(11.54) | 45(21.63) | 0.006 |
| | No | 184(88.46) | 163(78.37) | |
| Screening | Yes | 45(21.63) | 115(55.29) | <0.0001 |
| | No | 163(78.37) | 93(44.71) | |
| Sonography | Yes | 53(25.48) | 182(87.50) | <0.0001 |
| | No | 155(74.52) | 26(12.50) | |
| Blood Test | Yes | 109(52.40) | 183(87.98) | <0.0001 |
| | No | 99(47.60) | 25(12.02) | |
| First Child (sex) | Male (n=128) | 68(45.33) | 60(46.15) | 0.891 |
| | Female (n= 152) | 82(54.67) | 70(53.85) | |
| Second Child (sex) | Male (n= 58) | 32(43.84) | 26(55.32) | 0.219 |
| | Female (n= 62) | 41(56.16) | 21(44.68) | |
| Third Child (sex) | Male (n= 7) | 3(25.0) | 4(33.33) | 0.653 |
| | Female (n= 17) | 9(75.0) | 8(66.67) | |

Based on the results, the mean \pm standard deviation of pre-pregnancy mother body mass index (BMI)

was $27.83 \pm 4.18 \text{ kg/m}^2$ in the case group and $26.69 \pm 4.21 \text{ kg/m}^2$ in the control group. Other information is summarized in Table 2.

Table 2. Comparison of Demographic Information in Mothers with and without Abortion Based on One Sample t-Test.

| Variable | Group | n (%) | Mean \pm SD | Std Error | 95% CI | P-value |
|--|---------|-------------|-------------------|-----------|---------------|---------|
| Maternal age (year) | Control | 208 (100) | 31.00 \pm 5.52 | 0.38 | 30.25-31.76 | 0.038 |
| | Case | 208 (100) | 32.27 \pm 6.85 | 0.47 | 31.34-33.21 | |
| Husband age (year) | Control | 208 (100) | 35.25 \pm 5.95 | 0.41 | 34.44-36.06 | 0.065 |
| | Case | 208 (100) | 36.46 \pm 7.37 | 0.51 | 35.45-37.47 | |
| Weight (kg) | Control | 208 (100) | 70.79 \pm 11.74 | 0.81 | 69.18-72.39 | 0.260 |
| | Case | 208 (100) | 72.07 \pm 11.39 | 0.78 | 70.51-73.62 | |
| Height (cm) | Control | 208 (100) | 162.97 \pm 5.30 | 0.36 | 162.24-163.69 | 0.0001 |
| | Case | 208 (100) | 160.86 \pm 5.39 | 0.37 | 160.12-161.60 | |
| pre-pregnancy Body Mass Index ($\frac{\text{kg}}{\text{m}^2}$) | Control | 208 (100) | 26.69 \pm 4.21 | 0.29 | 26.11-27.27 | 0.005 |
| | Case | 208 (100) | 27.83 \pm 4.18 | 0.29 | 27.26-28.41 | |
| First Child (age-year) | Control | 130 (62.5) | 7.86 \pm 4.72 | 0.41 | 7.04-8.68 | 0.109 |
| | Case | 150 (72.11) | 8.88 \pm 5.73 | 0.46 | 7.96-9.81 | |
| Second Child (age-year) | Control | 47 (22.60) | 6.85 \pm 3.21 | 0.46 | 5.90-7.79 | 0.439 |
| | Case | 73 (35.10) | 7.45 \pm 4.63 | 0.54 | 6.37-8.53 | |
| Third Child (age-year) | Control | 12 (5.77) | 4.83 \pm 3.21 | 0.92 | 2.79-6.87 | 0.295 |
| | Case | 12 (5.77) | 6.91 \pm 5.91 | 1.70 | 3.15-10.67 | |

Results of logistic regression analysis showed that with the increase in mother's age, the risk for abortion increased (OR=1.03, 95% CI=1.00-1.06, p-value=0.039). After adjustment for other variables, with the increase in mother's age, the risk for abortion also increased significantly with an odds ratio of 4.67 (95% CI=1.11-19.62). Multiple regression logistic analysis confirmed a significant relationship between abortion and mother's age

(OR=4.67, 95% CI=1.11-19.62), previous cesarean delivery (OR=4.33, 95% CI=1.94-9.66), not taking folic acid pills during pregnancy (OR=14.92, 95% CI=6.25-35.61), low income (OR=7.00, 95% CI=2.02-24.35), not smoking (OR=0.28, 95% CI=0.08-0.94), and not suffering from pre-diabetes (OR=0.13, 95% CI=0.03-0.47) ($p < 0.05$). There were no significant associations between abortion and other studied risk factors (Table 3)

Table 3. Univariate and Multivariable Logistic Regression Analysis of Factors Associated with Abortion Among Pregnant Women.

| Variables | Category | Case group n (%) | Control group n (%) | Unadjusted OR (95% CI) | P-value | Adjusted OR (95% CI) | P-value |
|----------------------|-----------------------------------|------------------|---------------------|------------------------|---------|----------------------|---------|
| Mother age (year) | <40 | 181(87.02) | 202(97.12) | 1 | | 1 | |
| | >40 | 27(12.98) | 6(2.88) | 1.03 (1.00-1.06) | 0.039 | 4.67 (1.11-19.62) | 0.035 |
| Contraceptive method | Condom | 70 (33.81) | 107 (51.44) | 1 | | 1 | |
| | Natural family planning | 94 (45.41) | 57 (27.40) | 2.52 (1.61-3.93) | 0.001 | 4.46 (1.88-10.53) | 0.001 |
| | Combined pills | 25 (12.08) | 27 (12.98) | 1.41 (0.75-2.63) | 0.274 | 2.58 (0.89-7.42) | 0.078 |
| | Contraceptive injection | 9 (4.35) | 7 (3.37) | 1.96 (0.69-5.51) | 0.20 | 7.88 (1.18-52.48) | 0.033 |
| | IUD (intrauterine device or coil) | 9 (4.35) | 10 (4.81) | 1.37 (0.53-3.55) | 0.51 | 3.53 (0.71-17.43) | 0.120 |

| | | | | | | | |
|---|------------|-------------|-------------|---------------------|-------|---------------------|---------|
| Last delivery type | Vaginal | 89 (59.33) | 110 (84.62) | 1 | | 1 | |
| | Cesarean | 61 (40.67) | 20 (15.38) | 3.76 (2.11-6.71) | 0.000 | 4.33 (1.94-9.66) | <0.0001 |
| Acid folic pills (one month before pregnancy) | Yes | 96 (46.15) | 189 (90.87) | 1 | | 1 | |
| | No | 112 (53.85) | 19 (9.13) | 11.60 (6.73-20.01) | 0.000 | 14.92 (6.25-35.61) | <0.0001 |
| Maternal disorder | No | 191 (91.83) | 204 (98.08) | 1 | | 1 | |
| | Yes | 17 (8.17) | 4 (1.92) | 0.22 (0.07-0.66) | 0.007 | 3.07 (0.52-17.92) | 0.212 |
| Monthly income | ≤150 \$ | 42 (20.19) | 18 (8.65) | 1 | | 1 | |
| | 150-250 \$ | 112 (53.85) | 105 (50.49) | 3.67 (1.91-7.02) | 0.000 | 7 (2.02-24.25) | 0.002 |
| | ≥250 \$ | 54 (25.96) | 85 (40.86) | 1.67 (1.08-2.58) | 0.019 | 0.01 (0.46-2.21) | 0.965 |
| Smoking | No | 161 (77.40) | 194 (93.27) | 1 | | 1 | |
| | Yes | 47 (22.60) | 14 (6.73) | 0.24 (0.13-0.46) | 0.000 | 3.52 (1.05-11.81) | 0.041 |
| Pre-diabetes | No | 172 (82.69) | 200 (96.15) | 1 | | 1 | |
| | Yes | 36 (17.31) | 8 (3.85) | 0.19 (0.08-0.42) | 0.000 | 7.48 (2.10-26.54) | 0.002 |
| Mother blood group | O+ | 40 (19.23) | 96 (46.15) | 1 | | 1 | |
| | A+ | 106 (50.96) | 65 (31.25) | 3.91 (2.41-6.33) | 0.000 | 5.75 (2.43-13.57) | <0.0001 |
| | A- | 12 (5.77) | 1 (0.48) | 28.79 (3.62-228.93) | 0.001 | 10.71 (0.80-142.10) | 0.072 |
| | B+ | 30 (14.42) | 23 (11.06) | 3.13 (1.62-6.03) | 0.001 | 2.52 (0.69-9.23) | 0.16 |
| | B- | 1 (0.48) | 4 (1.92) | 0.6 (0.65-5.53) | 0.652 | 1.01 (0.02-43.50) | 0.99 |
| | AB+ | 16 (7.69) | 10 (4.81) | 3.84 (1.60-9.18) | 0.002 | 20.74 (4.52-95.12) | <0.0001 |
| | AB- | 0 | 1 (0.48) | 1 | | 1 | |
| | O- | 3 (1.45) | 8 (3.85) | 0.41 (0.28-0.60) | 0.000 | 2.40 (0.21-26.43) | 0.47 |

*OR= Odds Ratio, P≤0.05 significant meaning

Mothers under 30 years old who smoked exhibited a higher risk of abortion. The results of the interaction between maternal age and smoking are presented in Table 4. As maternal age increases, the correlation

between smoking and abortion risk significantly escalates, surpassing the individual effects of these variables. This phenomenon may be attributed to the cumulative impact of smoking over time.

Table 4. Multivariable Logistic Regression Analysis with the Interaction between Maternal Age and Smoking

| Variables | Category | Adjusted Odds Ratio | Std. Error | 95% CI | P-value |
|--|---------------------------|---------------------|------------|------------|---------|
| Interaction between Mother age and Smoking | <30 years and Not smoking | 1 | | | |
| | <30 years and smoking | 7.04 | 5.63 | 1.46-33.81 | 0.015 |
| | >30 years and Not smoking | 2.49 | 2.24 | 0.42-14.52 | 0.308 |
| | >30 years and smoking | 1.50 | 0.58 | 0.69-3.23 | 0.296 |
| Contraceptive Method | Natural family planning | 1 | | | |
| | Condom | 0.19 | 0.08 | 0.08-0.45 | <0.0001 |

| | | | | | |
|---|-------------------------|-------|-------|-------------|---------|
| | Combined pills | 0.58 | 0.30 | 0.20-1.62 | 0.302 |
| | Contraceptive injection | 1.47 | 1.42 | 0.22-9.84 | 0.688 |
| | IUD | 1.08 | 0.83 | 0.24-4.91 | 0.913 |
| Late delivery type | Vaginal | 1 | | | |
| | Cesarean | 5.21 | 2.10 | 2.36-11.51 | <0.0001 |
| Acid folic pills (one month before pregnancy) | Yes | 1 | | | |
| | No | 16.93 | 7.55 | 7.06-40.61 | <0.0001 |
| Pre-diabetes | Yes | 1 | | | |
| | No | 0.12 | 0.07 | 0.03-0.43 | 0.001 |
| Monthly income | ≤150 \$ | 1 | | | |
| | 150-250 \$ | 0.14 | 0.09 | 0.04-0.51 | 0.002 |
| | ≥ 250% | 0.15 | 0.09 | 0.04-0.53 | 0.003 |
| Mother blood group | O+ | 1 | | | |
| | O- | 2.02 | 2.61 | 0.15-25.60 | 0.587 |
| | A+ | 3.91 | 2.24 | 2.31-12.18 | <0.0001 |
| | A- | 11.64 | 15.26 | 0.89-152.11 | 0.061 |
| | B+ | 2.68 | 1.73 | 0.75-9.53 | 0.127 |
| | B- | 1.15 | 1.98 | 0.03-33.71 | 0.935 |
| | AB+ | 3.84 | 13.72 | 4.11-79.93 | <0.0001 |
| AB- | 1 | | | | |

Discussion

The current study revealed a significant increasing relationship between maternal age and abortion risk, with an adjusted odds ratio of 4.67, after adjusting for multiple covariates, which was statistically significant. Consequently, the risk of abortion in mothers with at least one gravidity was 8% less than nulliparous mothers. Therefore, mothers who had a history of more than two abortions significantly increased the risk of abortion.

The results of this study were consistent with previous studies regarding the association between advanced maternal age and abortion. It is observed that 75% of pregnancies occurred between 15-24 years old (16,17). Based on different studies, advanced maternal age will increase the risk of abortion due to fetal chromosomal disorders. It was reported that the higher risk of abortion came at a maternal age ≥45 years old (18). Erfani (2010) showed a high risk of abortion in maternal age over 35 years old (19). In another study conducted in the United States, maternal age was described as one of the abortion's risk factors (20).

It was reported that the highest abortion ratio was witnessed in women with diploma education, and

those who were housewives in both the case and control groups. This may be due to the majority of housewife mothers having a diploma. The risk of abortion was decreased in mothers with academic education (BSc and above), which may be attributed to the lower maternal ages.

The results of this study demonstrated that ABO blood type is associated with abortion. Most women (92.8%) and their husbands (77.6%) had Rh-positive blood groups. Mothers with blood type A negative had 7.35 times the risk of abortion compared to those with blood type A positive. Several epidemiological studies have reported the association between abortion and ABO blood types with Rh negativity (21). The higher representation of blood group type A was associated with recurrent abortion rather than ABO blood types (22). Other studies showed a relationship between pregnancy complications and blood group type A (23). Additionally, parental blood group incompatibility may increase the risk of abortion (24). However, the relationship between blood type and infertility and abortion is still controversial.

According to the results of this study, both direct and passive smoking significantly increased the risk

of abortion. Additionally, consuming more than 6 cups of tea or coffee per day was found to increase the risk of abortion by 2.77, which aligns with similar findings in the literature (25). Skogsdal (2022) reported that the odds of abortion increased with the number of cigarettes smoked per day, and there was also a significant relationship between abortion and caffeine or alcohol consumption (26). Previous reports suggest that the risk of abortion is 20-80% higher in addicted women (18,27), which may further elevate the risk of spontaneous abortion in nonsmoking women (28). Moreover, consuming 100 mg of caffeine increases the risk of spontaneous abortion by 22% (29).

In fact, caffeine found in coffee can elevate blood pressure and act as a diuretic, leading to dehydration, which can have negative effects on the body. If caffeine reaches the fetus during pregnancy, it can reduce blood flow to the placenta, potentially harming the fetus. Smoking during pregnancy is also known to increase the risk of health problems for both the mother and the fetus. Therefore, it is crucial to raise awareness among mothers about the negative impacts of smoking and excessive caffeine consumption during pregnancy, particularly in health centers.

According to the results of the present study, there was a significant relationship between maternal diseases (infections), a history of abnormal fetal birth, and abnormalities confirmed in amniocentesis with abortion. Additionally, a mother's history of pre-diabetes increased the likelihood of abortion. These findings are consistent with results from various other studies.

Bacterial infections rank among the most common infectious diseases worldwide, exerting adverse effects on sexual and reproductive health (30). Such infections can elevate the risk of ectopic pregnancy in women, leading to sudden and severe bleeding. Moreover, pregnant women afflicted with these infections may encounter complications such as miscarriage, preterm delivery, stillbirth, ectopic

pregnancy, and fetal abnormalities (31). Studies have indicated that maternal infection by mycoplasma is associated with abortion. Additionally, women with uncontrolled diabetes tend to experience a higher rate of miscarriage, although the precise mechanism underlying this phenomenon remains unknown (18).

In this study, a majority of 45.41% of mothers in the case group utilized natural family planning, while 51.44% opted for condom use in the control group as contraceptive methods, revealing a significant relationship. A study indicated that induced abortion was associated with mothers' use of natural family planning (32). Conversely, contraceptive pills served as a preventive factor, reducing the risk of abortion by 43%, a finding consistent with other studies (33,34). This might be attributed to the therapeutic effects of contraceptive pills, which, in addition to contraception, are sometimes employed to prevent ovarian cysts or strengthen follicles (35).

The results of this study demonstrated that the risk of abortion increased with the number of pregnancies, children, and cesarean deliveries, while children's sex did not emerge as a significant risk factor for abortion. These findings are in line with similar studies (36,37). One significant factor contributing to increased abortion risk is advanced maternal age, which heightens the likelihood of fetal chromosomal disorders (18). Additionally, the increased number of children, coupled with insufficient time for mothers to engage in physical activity and the influence of an unhealthy lifestyle, may elevate the likelihood of underlying diseases and abortion.

Furthermore, the risk of abortion was found to increase with decreasing monthly income. Prior research has demonstrated the impact of income on fertility behaviors (38). Mahmoudiani (2018) highlighted a higher incidence of abortion in lower-income families (37), and numerous studies have confirmed the correlation between lower income and abortion (39,40).

According to the results, mothers taking folic acid pills (one or three months before pregnancy) experienced a decreased risk of abortion, suggesting a preventive effect. A study conducted in England, examining the levels of riboflavin and folic acid in the serum of women experiencing spontaneous abortions in the first and second trimesters of pregnancy, revealed a high prevalence of deficiency, reaching 84% (41). This underscores the importance of screening pregnant women for nutritional deficiencies and recommending supplementation, such as folic acid, to prevent abortion and other pregnancy complications.

Future research

Future qualitative research studies could be done to define genetics factors resulting in abortion besides environmental factors among pregnant women.

Conclusion

Advanced maternal age and lower income were significantly associated with an increased risk of abortion. Additionally, suffering from pre-diabetes, a history of previous cesarean pregnancy, smoking, not taking folic acid pills, and maternal disorders (infection) were identified as potential risk factors for abortion. None of the covariates studied modified the association between maternal risk factors and abortion. Providing education, healthcare, and fostering continuous interaction between healthcare workers and mothers are essential measures to promote safe pregnancies and deliveries and decrease the risk of abortion.

Limitation and strengths of the study

In this study, the most significant limitation was the absence of information regarding potential early abortions that were either unreported by mothers or not recorded in the system. Additionally, the incompleteness of certain data in the electronic records of pregnant mothers or the outcomes of newborns posed another limitation to this project.

However, these limitations were mitigated to some extent through telephone or face-to-face follow-up efforts.

Acknowledgements

Researchers express their sincere gratitude to all mothers that participated in the study.

Financial support

This study was funded by Ilam University of Medical Sciences.

Conflict of interest

There are no conflicts of interest to declare.

Authors' contributions

HS and AK conceived the initial idea, collected data, and drafted the primary manuscript. SK assisted in the analysis of the data. DA, MM, and AH reviewed and provided critical feedback on the manuscript. All authors approved the final manuscript as submitted and agreed to be accountable for all aspects of the work.

References

1. Carp, H. J. A. Progestogens and pregnancy loss. *Climacteric*. 2018; 22:1-5. doi.org/10.1080/13697137.2018.1436166
2. Smorgick N, Mittler A, Ben-Ami I, Maymon R, Vaknin Z, Pansky M. Retained products of conception: What is the risk for recurrence on subsequent pregnancies? *Eur J Obstet Gynecol Reprod Biol*. 2018; 5(224):1-5. doi.org/10.1016/j.ejogrb.2018.03.006
3. Kaser D. The status of genetic screening in recurrent pregnancy loss. *Obstet Gynecol Clin North Am*. 2018; 45(1):143-154. DOI: 10.1016/j.ogc.2017.10.007
4. Ammon Avalos L, Galindo C, Li DK. A systematic review to calculate background miscarriage rates using life table analysis. *Birth Defects Res Part A Clin Mol Teratol*. 2012; 94(6):417-23. DOI: 10.1002/bdra.23014
5. Akhtari E. Pregnancy termination from ancient Persia till present time. *Iran J Obstet Gynecol Infertil*. 2017; 19(40):1-5. Doi: 10.22038/ijogi.2017.8442
6. Bearak J, Popinchalk A, Ganatra B, Moller A-B, Tunçalp Ö, Beavin C et al. Unintended pregnancy and abortion by income, region, and the legal status of abortion: estimates from a comprehensive model for 1990–2019. *Lancet Glob Health*. 2020 Sep; 8(9):e1152-e1161. doi: 10.1016/S2214-109X(20)30315-6.
7. Ganatra B, Gerds C, Rossier C, Ronald Johnson Jr B, Tunçalp O, Assifi A, et al. Global, regional and subregional classification of abortions by safety 2010-14: Estimates from a Bayesian hierarchical model. *Lancet*. 2017; 390(10110): 2372-81. DOI: 10.1016/S0140-6736(17)31794-4
8. Guttmacher Institute. Unintended pregnancy and abortion worldwide [Internet]. 2020. [updated 2020 July]. Available from: <https://www.guttmacher.org/factsheet/induced-abortion-worldwide>.
9. Rastegari A, Baneshi MR, Haji-Maghsoudi S, Nakhaee N, Eslami M, Malekafzali H, et al. Estimating the annual incidence of abortions in Iran applying a network scale-up approach. *Iran Red Crescent Med J*. 2014; 16(10): e15765. [In Persian]
10. Chan RL, Olshan AF, Savitz DA, Herring AH, Daniels JL, Peterson HB, et al. Severity and duration of nausea and vomiting symptoms in pregnancy and spontaneous abortion. *Human reproduction*. 2010; 25(11):2907-12. doi: 10.1093/humrep/deq260
11. Moradinazar Mehdi, Najafi Farid, Moradi Nazar Zeinab, Hamzeh Behrooz, Pasdar Yahya, Shakiba Ebrahim. Lifetime Prevalence of Abortion and Risk Factors in Women: Evidence from a Cohort Study. *Journal of Pregnancy*. 2020; 1-8. <https://doi.org/10.1155/2020/4871494>
12. Nazarpour S, Tehrani FR, Simbar M, Azizi F. Thyroid dysfunction and pregnancy outcomes. *Iranian journal of reproductive medicine*. 2015; 13(7):387-96. PMID: 26494985 PMID: PMC4609317
13. Maghsoudi R, Mirhosseini M. Spontaneous abortion and anti-thyroid antibodies in mother's serum. *Life Science Journal*. 2014; 11:41-4. <https://www.scopus.com/inward/record.uri?eid=2-s2....>
14. Pershad J, Mugerwa KY, Filippi V, et al. Prevalence and determinants of self-reported anxiety and stress among women with abortion-related complications admitted to health facilities in Eastern and Southern Africa: A cross-sectional survey. *Int J Gynecol Obstet*. 2022; 156 (1): 53–62. doi:10.1002/ijgo.14042
15. Frederico M, Michielsen K, Arnaldo C, Decat P. Factors influencing abortion decision-making processes among young women. *Int J Environ Res Public Health*. 2018;15(2):329-42. doi: 10.3390/ijerph15020329
16. Alexander P, Frick MD MRCOG. Advanced maternal age and adverse pregnancy outcomes. *Journal of Best Practice & Research Clinical Obstetrics & Gynaecology*. 2021: 92-100. <https://doi.org/10.1016/j.bpobgyn.2020.07.005>
17. Rosaly Correa-de-Araujo, Sung Sug (Sarah) Yoon. Clinical Outcomes in High-Risk Pregnancies Due to Advanced Maternal Age. *Journal of Women's Health*. 2021: 30(2). <https://doi.org/10.1089/jwh.2020.8860>
18. The practice committee of the American society for reproductive medicine. Evaluation and treatment of recurrent pregnancy loss: a committee opinion. *Fertility and Sterility*. 2012; 98 (5): 1103-11.
19. Erfani A. Tehran survey of fertility. A study on the reproductive behavior of married women in Tehran. Tehran: The population studies and research center in Asia and pacific (PSRC). 2010. <https://www.scirp.org/reference/referencespapers?referenceid=1170785>
20. Martin JA, Hamilton BE, Osterman MJK, Hyattsville, MD. Births: Final data for 2016. National vital statistics reports. sNational Center for Health Statistics, 2018: 67(1).
21. Pourjafari H, Hashemzadeh Chaleshtori M, Arab M. Frequencies of antigens and their alleles from ABO & RH blood types in a group of women with two or more abortions. *Avicenna Journal of Clinical Medicine* 2004; 10(4):43-46. <https://www.sid.ir/paper/538877/en>
22. Shahverdi E, Moghaddam M, Talebian A, Abolghasemi H. Distribution of blood groups in the Iranian general population. *J Blood Group Serol Mol Genet*. 2016; 32(4):135. PMID: 28257227
23. Higgins JR, Walshe JJ, Darling MR, Norris L, Bonnar J. Hemostasis in the uteroplacental and peripheral circulations in normotensive and pre-eclamptic pregnancies. *Am J Obstet Gynecol*. 1998; 179(2): 520-6. DOI: 10.1016/s0002-9378(98)70389-8
24. Hassanzadeh-Nazarabadi M, Shekouhi S, Seif N. The Incidence of Spontaneous Abortion in Mothers with Blood Group O Compared with other Blood Types. *Int J Mol Cell Med*. 2012; 1(2):99-104. PMID: 24551765 PMID: PMC3920496
25. Karbasian Zahra. The relationship between pregnant mothers addiction with abortion. *New advances in psychology, education, and education*. 2018; 1(2): 59-67.
26. Skogsdal Y, Karlsson J, Tydén T, Patil S, Backman H. The association of smoking, use of snuff, and preconception alcohol consumption with spontaneous abortion: A population-based cohort study. *Acta Obstet Gynecol Scand*. 2023;102:15-24. doi: 10.1111/aogs.14470
27. Triggianese P, Perricone C, Perricone R, De Carolis C. Prolactin and Natural Killer Cells: Evaluating the

- Neuroendocrine-immune Axis in Women with Primary Infertility and Recurrent Spontaneous Abortion. *Am J Reprod Immunol*. 2014. DOI:10.1111/aji.12335
28. Khazaei S, Soltani O. Caffeine intake and spontaneous abortion. *SCIENTIFIC JOURNAL of Ilam Medical University*. 2003; 10(36, 37): 39-45. DOI: 10.1056/NEJM200012213432503
 29. Kitts DD. Caffeine metabolism and its deposition in the pregnant rat and fetus. *Can Inst Food Sci Technol J* 1986 ; 19: XL.
 30. Pustejovsky JE, Rodgers MA. Testing for funnel plot asymmetry of standardized mean differences. *Res Synth Methods*. 2019; 10: 57–71. doi.org/10.1002/jrsm.1332
 31. Afolabi BB, Moses OE, Oduyebo OO. Bacterial vaginosis and pregnancy outcome in Lagos, Nigeria. *Open Forum Infect Dis*. 2016; 3:ofw030. doi:10.1093/ofid/ofw030.
 32. Erfani A. Induced abortion in Tehran, Iran: Estimated rates and correlates. *Int Perspect Sex Report Health*. 2011; 37(3): 134-42. DOI: 10.1363/3713411
 33. Ford J.H, MacCormac L. Pregnancy and lifestyle study: the long-term use of the contraceptive pill and the risk of age-related miscarriage. *Human Reproduction*. 1995; 10(6): 1397–1402. DOI: 10.1093/humrep/10.6.1397
 34. L. Michie, S.T. Cameron. Emergency contraception and impact on abortion rates. *Journal of Best Practice & Research Clinical Obstetrics & Gynaecology*. 2020: 111-119. <https://doi.org/10.1016/j.bpobgyn.2019.06.008>
 35. Kovacs P, Barg P.E, Witt B.R. "Hypothalamic-pituitary suppression with oral contraceptive pills does not improve outcome in poor responder patients undergoing in vitro fertilization-embryo transfer cycles," *Journal of Assisted Reproduction and Genetics*. 2001; 18(7): 391–394. DOI: 10.1023/a:1016626607387
 36. Hansen M-LH, Mølgaard-Nielsen D, Knudsen LB, Keiding N. Rates of induced abortion in Denmark according to age, previous births and previous abortions. *Demographic Research*. 2009; 21: 647-80. Doi: 10.4054/DemRes.2009.21.22
 37. Mahmoudiani S, Ahmadi AY, Javadi A. The Prevalence and Influential Factors of Abortion in the Women in the Rural Areas of Fars Province, Iran (2015). *Iran Journal of Nursing*. 2018; 31(115):51-61. Doi: 10.29252/ijn.31.115.51
 38. Abbasi Shavazi Mj, Khani S. Economic Insecurity, Marriage and Fertility Ideals: A study among Mothers and Children Generation in Sanandaj District, *Journal of Population Studies*. 2016; 1(2): 63-99. Doi: 20.1001.1.1735000.1393.9.17.2.8
 39. Alemayehu M, Yebyo H, Medhanyie AA, Bayray A, Fantahun M, Goba GK. Determinants of repeated abortion among women of reproductive age attending health facilities in Northern Ethiopia: a case-control study. *BMC Public Health*. 2017; 17(1): 188-96. DOI: 10.1186/s12889-017-4106-1
 40. Chinichian M, Halakooii-Naiini K, Rafaii-Shirpak K.. Voluntary Abortion in Iran: a qualitative study, *Journal of Payesh*. 2007; 6(3): 219-32. URL: <http://payeshjournal.ir/article-1-701-en.html>
 41. Neela J, Raman L. The relationship between maternal nutritional status and spontaneous abortion. *Natl Med J India*. 1997; 10(1):15-6. PMID: 9069700.