# Cucurbita maxima (Pumpkin) seeds: Scolicidal activity and preventive efficacy of its extract on experimental hydatidosis in mice

Arash Babaei\*1, Arash Jafari2, Mohammad Asadpour2, Morteza Shamsi3

- 1. Department of Biology, Faculty of Sciences, Malayer University, Malayer, Iran
- 2. Department of Pathobiology, Faculty of Veterinary Medicine, Shiraz University, Shiraz, Iran
- 3. Department of Parasitology, Faculty of Paramedical Sciences, Ilam University of Medical Sciences, Ilam, Iran

\*Corresponding author:Tel: +98 8133339942 Fax: +98 8133339942

Address: Department of Biology, Faculty of Sciences, Malayer University, Malayer, Iran

E-mail: a.babaei@sheffield.ac.uk

Received; 2017/06/13 revised; 2017/07/5 accepted; 2017/08/21

#### Abstract

**Introduction:** Hydatid Disease (HD), a helminth infection with various clinical complications is a serious economic and public health concern around the world. Nowadays, different agents have been broadly applied as scolicids; these agents have several side effects on patients. Therefore, the aims of this experimental study were to evaluate the scolicidal activity of Cucurbita maxima seeds methanolic extract (ME) against protoscolices of hydatid cyst in-vitro. Also, an in vivo assay was designed to prove the preventive effects of the extract.

**Materials and methods:** For this work, Two different concentrations (25 and 50 mg/ml) of the C. maxima seeds ME were used in various exposure times (10, 20, 30, 40, 50, and 60 min). Additionally, thirty mice were infected intraperitoneally by injection of 1000 viable protoscolices and were divided into three groups with ten mice in each Group I (treated by Albendazole), group II (received 50 mg/ml of C. maxima seeds ME) and group III (untreated controls).

**Results:** Overall, C. maxima seeds ME exhibited significant scolicidal effects in compared with control group (P<0.05). At a concentration of 50 mg/ml, the mortality percentage increased to 84.7%, 96.1% and 100% after 40, 50 and 60 minutes, respectively. Also, the obtain data in necropsy demonstrated that C. maxima seeds ME have an inhibitory effect on the hydatid cyst development. No hydatid cyst formation observed in mice treated with C. maxima seeds ME (50 mg/ml) versus control group which hydatid cyst occurred in 6 (from 10) mice.

**Conclusion:** The current investigation indicated that applying C. maxima seeds ME could be considered as a potential scolicidal agent due to being economical, safer and non-toxic compared to the reference chemical drugs. However, further studies on probably active compounds and also to the determination of its mode of action on parasites are required.

Keywords: Hydatid disease, Cucurbita maxima methanolic extracts, Scolicidal, Preventive assay

#### Introduction

Hydatid Disease (HD) is a major neglected zoonotic disease with various clinical complications caused by the larval stage of Echinococcus granulosus (1, 2). HD is still an important public health concern in many countries of the world, such as Iran.

**Copyright © 2018 Journal of Basic Research in Medical Science.** This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>) which permits copy and redistribute the material, in any medium or format, provided the original work is properly cited.

The disease affects humans as well as domestic animals, including cattle, sheep, camels, pigs, horses and others worldwide (3, 4). HD affects mainly the liver (50-70% of all cysts) but can develop in lung (20-30%) and, less frequently, in spleen, bone, brain, and other organs (5, 6). The preferred method for treatment of HD is surgery, but it increases the risk of intraoperative spillage of protoscolices (6,7,8). This is the major cause of recurrence, which is seen in approximately 10% of the postoperative cases. Operative spillage also lead to the secondary disseminated intraperitoneal hydatidosis (6, 8). There is a severe need to some safe and more effective scolicids for killing or inactivation of protoscolices in surgeries (7, 9). Formalin, hypertonic saline, cetrimide, chlorhexidine, hydrogen peroxide and ethyl alcohol are some of the compounds used as scolicidal, but all of these agents are concentration dependent and their degree of dilution in the cyst contents is quite unpredictable (10), Also, they have different adverse effects such as liver necrosis and biliary tract fibrosis (2). Thus, it is indispensable to develop new scolicidal materials with more efficacious and low adverse effects, which can be during implemented surgery Considerable evidence has demonstrated that some plant derived products (PDPs) can be useful as scolicidal (12). Based on previous in vitro and in vivo data on scolicidal effects of PDPs, **Mallotus** philippinensis (13),Allium sativum Zingiber (Garlic) (6), (14),Zataria multiflora (11) and Berberis vulgaris (15) showed significant scolicidal activity. Cucurbita maxima (Pumpkin) (Family commonly Cucurbitaceae) known Squash is widely used as vegetable and a source of vitamin A, iron, phosphorus and calcium (16). C. maxima seeds with considerable safety margin versus commonly consumed chemical drugs (17), is a very rich in terms of amino acids that are building blocks of proteins, especially amino essential acids such

phenylalanine and methionine which are not produced in the human body(18). Investigation on C. maxima (Squash) reported that spinasterol isolated from the flowers of C. maxima potentially showed anti-carcinogenic, anti-genotoxic (19), and anti-mutagenic activity (20). The C. maxima seeds were used in the treatment of digestive disorders [21], as well as essential oil of C. maxima seeds exhibited significant anti helmintic activity (16, 18, 22). Recently, an in vitro study on hydatid cyst protoscolices revealed that 50 mg/ml maxima (pumpkin) C. chloroformic extracts showed scolicidal on the previous activity (7). Based investigations on anthelmintic scolicidal activity of C. maxima seed extracts, present study was conducted in two stages, first was the in vitro activity of C. maxima methanolic extracts (ME) on protoscolices of Hydatid Cyst, and also in the next stage in vivo efficacy of this methanolic compound for the prevention hydatid cysts tested mice experimentally infected with protoscolices of hydatid cyst.


## Materials and methods

Collection of protoscolices: Protoscolices were collected from livers of naturally infected sheep, with hydatid disease, slaughtered at Malayer abattoir, Hamadan Southwestern province, of Iran. Protoscolices was isolated from the cysts ,according to the method described previously by Moazeni and Nazer (6). Briefly, the fluid of cysts was transferred cylinders by aspiration, aseptically and left to set for 30 min. The protoscolices settled down at the bottom. The supernatant was discarded and the remaining protoscolices at the bottom were washed three times with normal saline. The viability of the protoscolices was tested by their motility using a light microscope. The protoscolices transferred into a dark container containing normal saline solution and stored at 4°C until use.

**Methanolic extract Preparation:** The *C*. maxima seeds were purchased from a local market in Hamadan (Iran) and C. maxima seeds ME was obtained as follows: C. maxima seeds, dried under shade, and mechanically powdered using commercial electric blender. To obtain the methanolic extract, 200 g of dry seeds powder was added to 800 ml of pure methanol and mixed gently for 2h using a magnetic stirrer. The obtained solution was left at room temperature for 24h. The compound was stirred again and filtered and then the solvent was removed by evaporation in a rotating evaporator. The remaining semisolid material was then freeze-dried. The obtained residue was placed into a glass container and stored at 4°C until use (6, 23).

**Scolicidal activity:** In the present study, we had two concentrations (25 and 50 mg/ml) of the *C. maxima* seeds ME for 10, 20, 30, 40, 50, and 60 min. To obtain the *C.maxima* seeds ME solution at 25 and 50 mg/ml concentrations, 0.25 and 0.5 gr of dried extract was dissolved in 10 ml of normal saline, respectively. Then 2 ml of

C. maxima seeds ME solution was placed test tubes, to which drop off protoscolex-rich sediment was added. The contents of the tubes were mixed, gently. The tubes were incubated at  $37^{\circ C}$  for 10, 20, 30, 40, 50, and 60 min. At the end of each incubation time the upper phase was carefully discarded without disturb the protoscolices. To check the viability of protoscolices, 0.1% Eosin stain (1g of Eosin powder in 1000 ml distilled water) was added to the settled protoscolices and mixed gently. The upper portion of the solution was discarded after 15 min of incubation. The remaining protoscolices was then smeared on a manually scaled glass slide, covered with a cover glass, and examined under a light microscope. The percentages of dead protoscolices (which absorbed Eosin and stained red after 15 min of incubation) were calculated by counting a minimum of 500 protoscolices. Remained colorless protoscolices were considered potentially Non-treated viable (Figure 1). protoscolices were considered as a control experiment. in each experiments were performed in triplicate (6, 7).



**Figure 1.** (Left) Live protoscolices after staining with 0.1% eosin. (Right) Dead protoscolices after exposure to *C.maxima* seeds ME and staining with 0.1% eosin.

**Experimental infection of mice:** Preventive trial in the present study was designed according to the method described previously (11) with some modification as follows: To confirm the

preventive effects of *C. maxima* seeds ME on hydatid cyst formation, an in vivo assay was carried out on 30 healthy laboratory mice, weighing 20 to 30 g, 6-8 weeks old.

Field application and study design: The mice were infected intraperitoneally by injection of 1000 protoescolices per animal, dissolved in 0.5 ml of distilled water. They were randomly divided into three groups of ten animals each. The experimental groups were kept in cages separately with free access to food and water. Group I (10 mice) was treated by Albendazole 150 mg/kg at body weight/day for 10 days, the group II (10 mice) received 50 mg/ml of C.maxima seeds ME in drinking water for 6 months, and group III (10 mice) were the untreated controls. Albendazole and C. maxima seeds ME each was administered by oral gavage. All mice (from groups I, II and III were euthanized at the end the experiment (after 6 months) by Ether, and a necropsy was carried out immediately. At necropsy, the body cavity was opened, and the internal organs were observed for hydatid cysts (11).


# Statistical analysis

The scolicidal trial was repeated three times and their average of death protoscolices calculated were and considered for each extract. Statistical analysis was performed using SPSS software version 20. Differences between the control and treatment groups were analyzed by Chi-square test. Values were statistically considered significant at p < 0.05.

#### Results

Scolicidal effects of C. maxima seeds ME: The mortality rate of hydatid cyst protoscolices after exposure to different concentrations of the C. maxima seeds ME following various exposure times are presented in (Table 1). It could be observed that the C. maxima seeds ME in mg/ml concentrations exhibited significant scolicidal effects in compared with control group (P<0.05). All protoscoleces were killed after 60 min of

exposure to 50 mg/ml concentration of the agent. Also, after 60 min exposure time, the scolicidal activity of C. maxima seeds ME at a concentration of 25 mg/ml was The results showed a time dependent manner in mortality rate of hydatid cyst protoscolices, so that with the increasing in time mortality percentage was increased, dramatically (Figure 2). When protoscolices were exposed to the *C*. maxima seeds ME at a concentration of 50 mg/ml, the mortality percentage increased to 84.7%, 96.1% and 100% after 40, 50 60 minutes. respectively. mortality rate of hydatid cyst protoscolices after exposure to 25 mg/ml concentration of C. maxima seeds ME at all times was lower versus 50 mg/ml of its. The results of our study confirmed that C. maxima seeds ME showed a high scolicidal activity in vitro.



**Figure 2.** Mortality percentage of protoscolices in different concentrations of the C.maxima seeds ME and various times. Values are presented as Mean of mortality rate (%) of protoscolices.

**Preventive effects of** *C. maxima* **seeds ME:** We used 50 mg/ml concentration of *C. maxima* seeds ME in Preventive trial, which exhibited higher scolicidal activity in vitro. Based on necropsy evidences, *C. maxima* seeds ME showed a preventive effect on formation of hydatid cyst in mice. The obtain data in necropsy showed that no hydatid cyst formation observed in mice treated with *C. maxima* seeds ME (50 mg/ml in drinking water for 6 months) and Albendazole (150 mg/kg body weight/day for 10 days). Additionally, hydatid cyst

occurred in 6 (from 10) mice of control group, and the number of developed cysts

in this group were more than four in two mice.

Table1. Scolicidal activity of C.maxima seeds ME in different concentrations and various exposure times.

		Dead protoscoleces /total treated protoscoleces (%)					
Concentration	Experiment	10 min	20 min	30 min	40 min	50 min	60 min
25 mg/ml	1	127/500(21.9)	117/420(27.8)	255/560(45.6)	280/480(58.4)	387/500(77)	462/550(84)
	2	100/500(20)	152/540(28.2)	189/510(37)	258/520(49.6)	295/490(60)	384/460(83.4)
	3	117/520(22.5)	95/340(28)	176/480(36.7)	235/460(51)	305/390(78)	324/390(83)
Total		344/1600(21.5)	364/1300(28)	620/1550(40)	773/1460(53)	987/1380(71.5)	1170/1400(83.4)
50 mg/ml	1	182/480(37.9)	368/600(61.3)	434/600(72.3)	323/380(85)	412/430(96)	590/590(100)*
_	2	201/540(37.2)	252/420(60)	294/420(70)	522/600(87)	472/500(94.5)	450/450(100)*
	3	160/460(34.7)	281/480(58.5)	403/560(72)	443540(82)	518/530(97.8)	600/600(100)*
Total		543/1480(36.6)	901/1500(60)	1131/1580(71.5)	1288/1520(84.7)	1402/1460(96.1)	1640/1640(100)
Control	1	23/520(4.5)	38/580(6.5)	46/500(9.2)	46/380(12)	74/510(14.5)	84/600(14)
	2	26/480(5.4)	22/420(5.2)	23/290(8)	45/450(10)	63/460(13.7)	61/510(12)
	3	13/370(3.5)	27/360(7.5)	47470(10)	64/490(13)	91/600(15.2)	54/360(15)
Total		62/1370(4.4)	87/1360(6.4)	116/1260(9)	155/1320(11.7)	228/1570(14.5)	199/1470(13.6)

Values are presented as a fraction or percent of killed protoscolices.

# **Discussion**

Hydatid cyst larval stage as a (metacestode) of E. granulosus is still a serious economic issue and public health concern worldwide. Few strategies are available for the treatment of hydatid disease in human (3). Surgery is the most common method for particular WHO stage Chemotherapy disease. benzimidazoles and also PAIR (puncture, aspiration, injection and reaspiration) are recommended as alternative strategies to surgery, in some patients who cannot tolerate surgery (13). Removal of the cysts with chemotherapy using benzimidazoles before and after surgery is the best and selective method. But, there are evidences of resistance to synthetic anthelmintic agents and the adversity of numerous side thrombocytopenia, effects, including leukopenia and hepatotoxicity (4, 24). Therefore, investigations of appropriate alternative and safer compounds with a different mode of action to develop new treatment strategies are necessary. Several experimental studies conducted for finding such scolicidal, previously. For instance, hypertonic saline (25), silver nitrate (26), and chlorhexidine gluconate (27) were tested. Recently, natural compounds such PDPs were introduced good candidates to safer control agents that may provide scolicidal activity and

drawbacks. A number of researches describe the scolicidal effects of different herbs and spices and their volatile components. Recently, Moazeni et al. in some in vitro investigations, reported that herbal extracts including Sumac (Rhus coriaria L.) (28), Zingiber officinale R (14). Have a significant scolicidal activity. C. maxima seeds (Pumpkin) (Family Cucurbitaceae) which commonly known as Squash has been used in different parts of the world as traditional medicine for treatments of gastrointestinal parasites as anthelmintic. Also, anthelmintic and in vitro scolicidal activity of C. maxima seeds was confirmed, recently (7, 18, 29). This work for the first time describes the scolicidal effects of C. maxima seeds ME against protoscoleces of hydatid cysts on in vitro model. Additionally, preventive effects of C. maxima seeds ME on hydatid cysts was evaluated in a mouse model. Our results showed that C. maxima seeds ME potentially have a scolicidal activity, especially at concentrations 25 and 50 mg/ml (83.4 and 100% mortality rate) after 60 min of application, respectively. Scolicidal effects of C. maxima seeds ME against protoscoleces of hydatid cysts in previous studies which conducted by Eskandarian (7) was lower than present data, it could be due to differences in the

<sup>\*</sup>Significant difference at P<0.05.

extraction methods and concentrations. In addition to the confirmation of the Scolicidal effects of C. maxima seeds ME, especially in the present study, we also confirmed our data in vivo assay. Based on in vivo data no hydatid cyst formation observed in mice treated with C. maxima seeds ME (50 mg/ml in drinking water for 6 months) versus control group. Our experimental study proved the preventive effects of C. maxima seeds ME for hydatid cyst development. Although spinasterol is an active compound which isolated from flowers of C. maxima [19], further studies on probable other active compounds of C. maxima seeds ME to the determination of its mode of action on parasites are needed. demonstrated that C. Obviously, we maxima seeds ME (at 50 mg/ml concentration) exhibited a highly

#### References

- 1. Sarvi S, Daryani A, Sharif M, Rahimi MT, Azami D, Marhaba Z, et al. Domestic dog as a human health hazard in north of Iran. J Parasit Dis. 2014;2(4): 1-5.
- 2. Rahimi MT, Ahmadpour E, Rahimi-Esboei B, Spotin A, Kohansal- Koshki MH, Alizadeh A, et al. Scolicidal activity biosynthesized of silver nanoparticles against E.granulosus protoscolices. Int J Surg. 2015; 19(1):128-33.
- 3. Eckert J, Deplazes P. Biological epidemiological and clinical aspects of echinococcosis a zoonosis of increasing concern. Clin Microbiol Rev. 2004;17(3):107-35.
- 4. Mahmoudvand H, Fasihi-Harandi M, Shakibaie M, Aflatoonian MR, ZiaAli N, Sadat-Makki M, Jahanbakhsh S. Scolicidal effects of biogenic selenium nanoparticles against protoscolices of hydatid cysts. Int J Surg 2014; 12(4): 399-403.
- 5. Ammann RW, Eckert J. Cestodes echinococcus. Gastroenterol Clin North Am. 1996; 25(6):655-89.

significant scolicidal and preventive activity in vitro and in vivo trials, respectively. Based on our present data *C. maxima* seeds ME with considerable safety margin versus chemical drugs commonly consumed to treatment of HD (17), could be a logical candidate to use as an effective scolicids and preventive agent against cystic echinococcosis.

## **Conclusion**

The current investigation indicated that applying C. maxima seeds ME could be considered as a potential scolicidal agent due to being economical, safer and nontoxic compared to the reference chemical drugs. However, further studies on probably active compounds and also to the determination of its mode of action on parasites are required.

- 6. Moazeni M, Nazer A. In vitro effectiveness of garlic Allium sativum extract on scolices of hydatid cyst. World J Surg. 2010; 34(2): 2677-81.
- 7. Eskandarian, AA. Scolicidal effects of squash Corylus spp seeds, hazel Curcurbia spp nut and garlic Allium sativum extracts on hydatid cyst protoscolices. J Res Isfahan Uni Med Sci. 2012; 17(3): 1011.
- 8. Yang YR, Sun T, Li Z. Community surveys and risk factor analysis of human alveolar and cystic echinococcosis in Ningxia Hui autonomous region China. Bull World Health Org. 2006; 84(3): 714-21.
- 9. Paksoy Y, Odev K, Sahin M, Arslan A, Koc O. Percutaneous treatment of liver hydatid cysts comparison of direct injection of albendazole and hypertonic saline solution. Am J Roentgenol. 2005; 185(6):727-34.
- 10. Besim H, Karayalcin K, Hamamci O, Gungor C, Korkmaz A. Scolicidal agents in hydatid cyst surgery. HPB Surg. 1998; 10(1):347-51.

- 11. Moazeni, M, Larki S, Saharkhiz MJ, Oryan A, Ansary Lari M, Mootabi Alavi A. In vivo study of the efficacy of the aromatic water of Zataria multiflora on hydatid cysts. Antimicrob Agents Chemother. 2014; 58(3): 6003-8.
- 12. Gessler M, Nkunya MHH, Mwasumbi LB, Heinrich M, Tanner M. Screening Tanzanian medicinal plants for antimalarial activity. Acta Trop. 1994; 56(1): 65-77.
- 13. Gangwar M, Verma VC, Singh TD, Singh SK, Goel RK, Nath G. In vitro scolicidal activity of Mallotus philippinensis Muell arg fruit glandular hair extract against hydatid cyst E.granulosus. Asian Pac J Trop Med. 2013; 6(4): 595-601.
- 14. Moazeni M, Nazer A. In vitro lethal effect of Zingiber officinale R. on protoscolices of hydatid cyst from sheep liver. Microbiol Res. 2011; 2(2): 25.
- 15. Rouhani, S, Salehi N, Kamalinejad M, Zayeri F. Efficacy of Berberis vulgaris aqueous extract on viability of E. granulosus protoscolices. J Invest Surg. 2013; 26(2): 347-51.
- 16. Sengupta R, Banik J. Comparative studies on anthelmintic potential of C. maxima (pumpkin) seeds and Carica papaya seeds. Int J Res Ayurveda Pharm. 2013; 4(4):223-6.
- 17. Cruz R, Meurer CD, Silva E, Schaefer C, Santos A, Cruz B, Cechinel V. Toxicity evaluation of C. maxima seed extract in Mice. Pharm Biol. 2006; 44(2): 301-3.
- 18. Ayaz, E, Gokbulut C, Coskun H, Turker A, Ozsoy S, Ceylan K. Evaluation of the anthelmintic activity of pumpkin seeds in Mice naturally infected with Aspiculuris tetraptera. J Pharmacognosy Phytother. 2015; 7(9): 189-93.
- 19. Villasenor IM, Lemon P, Palileo A, Bremner JB. Antigenotoxic spinasterol

- from C.maxima flowers. Mutat Res Environ Mutagen. 1996; 360(2): 89-93.
- 20. Villaseñor IM, Domingo AP. Anticarcinogenicity potential of spinasterol isolated from squash flowers. Teratog Carcinog Mutagen. 2000; 20(3): 99-105.
- 21. Prokhvatilova S, Mikhalev V, Prokhovora LV, Evtushenko NS, Antonova NP, Simonova EP. HPLC determination of α tocopherol in tykveol preparation. Farmatsiya Moscow. 1998; 47(5): 31-2.
- 22. Ahmet-Basaran A, Faruk-Ciftci K, Kuesmenoglu S. Characteristics of Turkish C. maxima Duch seed oil. Acta Pharm Turc. 1998; 40: 17-20.
- 23. Razavi SM, Asadpour M, Jafari A, Malekpour SH. The field efficacy of Lepidium latifolium and Zataria multiflora methanolic extracts against Varroa destructor. Parasitol Res. 2015; 114(11): 4233-8.
- 24. Smego RA, Sebanego P. Treatment options for hepatic cystic echinococcosis. Int J Infect Dis. 2005; 9(2): 69-76.
- 25. Kayaalp C, Balkan M, Aydin C, Ozgurtas T, Tanyuksel M, Kirimlioglu V, et al. Hypertonic saline in hydatid disease. World J Surg. 2001; 25(8): 975-9.
- 26. Rajabi MA. Fatal reactions and methaemoglobinaemia after silver nitrate irrigation of hydatid cyst. Surg Pract. 2009; 13(1): 2-7.
- 27. Topcu O, Sumer Z, Tuncer E, Aydin C, Koyuncu A. Efficacy of chlorhexidine gluconate during surgery for hydatid cyst. World J Surg. 2009; 33(6): 1274-80.
- 28. Moazeni M, Mohseni M. Sumac Rhus coriaria scolicidal activity on hydatid cyst protoscolices. Surg Sci. 2012; 3(9): 452.
- 29. Srivastava M, Singh S. Anthelmintic activity of C. maxima Kaddu seeds. Indian J Med Res. 1967; 55(6): 629-32.