

Association of ESR- α and ESR- β gene polymorphisms with implantation failure in IVF-treated women in northwest of Iran: A case-control study

Parisa Pagard¹, Seyed Ali Rahmani^{2*}, Lida Heidari³

1. Department of Genetic, Tabriz Branch, Islamic Azad University, Tabriz, Iran
2. Department of Medical Genetics, Tabriz University of Medical Sciences, Tabriz, Iran
3. Gynecology Department, Eastern Azerbaijan ACECR ART Center, Eastern Azerbaijan Branch of ACECR, Tabriz, Iran

*Corresponding author: Tel: +98 9121961050 Fax:

Address: Department of Medical Genetics, Tabriz University of Medical Sciences, Tabriz, Iran

E-mail: rahmaniseyedali@yahoo.com

Received; 15/04/2020 Revised; 26/06/2020 Accepted; 11/07/2020

Abstract

Introduction: Estrogen, a crucial hormone during pregnancy, acts through two types of receptors. The estrogen receptor alpha and beta (ESR- α and ESR- β) are more abundant and exists in all human reproductive systems. Association of ESR- α and ESR- β genes polymorphisms has been reported in some reproductive problems such as spontaneous abortion, endometriosis-related infertility, and *in vitro* fertilization failure. In the present study, we investigated association between single nucleotide polymorphisms rs9340799 and rs2234693 (ESR- α) and rs1256049 and rs4986938 (ESR- β) with implantation failure in Iranian women.

Materials and Methods: In this case-control study, we collected 60 women with implantation failure as case group and 60 age and ethnically matched IVF-treated women with successful implantation as controls. Extraction of genomic DNA of both case and control members was performed using salting out method. The case and control groups genotyping was performed using tetra-primer amplification refractory mutation system-polymerase chain reaction (Tetra-ARMS PCR) method.

Results: There were no significant differences in the frequencies of genotype and allele frequency in ESR- β gene rs4986938 polymorphism between patients and control groups ($p>.005$). In contrast, we observed a significant difference in the frequencies alleles and genotypes of rs9340799 and rs2234693 (ESR- α) and rs1256049 (ESR- β) polymorphisms between patients and control groups.

Conclusion: We demonstrated that rs9340799 and rs2234693 (ESR- α), and rs1256049 (ESR- β) polymorphisms may play important role in implantation failure in women in northwest of Iran. However, more studies on different geographic areas, races and ethnicities are required to determine exact role of ESR- α and ESR- β genes polymorphisms in implantation failure.

Keywords: Polymorphisms, ESR- α , ESR- β , Implantation failure

Introduction

During implantation process, a receptive uterus is an essential factor that guaranties success of implantation and pregnancy (1). Factors, involved in the process, prepare the uterine luminal epithelium to its best

situation for blastocyst reception that finally makes a cross-link between uterus and blastocyst (2). Here, estrogen hormone plays a crucial role throughout the entire pregnancy, from regulating the production of progesterone to fetal development. This important role of estrogen has been proven in

Copyright © 2020 Journal of Basic Research in Medical Science. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits copy and redistribute the material, in any medium or format, provided that the original work is properly cited.

the uterine preparation for the implantation in mice (3).

Estrogen action is observed both on peripheral and central nervous system which is mediated by estrogen receptors (ER). Estrogen receptors α (ESR α , ESR1) and β (ESR β , ESR2) genes are involved in controlling the physiological response to estrogen (4). ESR α gene is present on chromosome 6q25.1-q25.2. It consists of 8 exons which are parted by 7 intronic regions having a complete size of 140 kb. The ESR β gene is located on chromosome 14q23.2-q23.3, comprising 8 exons and covers approximately 40 kb (5). ESR α codes for a 595 amino acid protein, while ESR β codes for a 530 amino acid protein (6).

Most widely studied polymorphisms of ESR α gene is rs2234693 and rs9340799, and ESR β gene is rs1256049 and rs4986938. The rs2234693 polymorphism occurs due to T/C transition in first intron, while the rs9340799 polymorphism is caused by a G/A transition located 50 base pairs downstream of the polymorphic site (7). Several sequence variants of the ESR β gene have also been identified which include 2 silent polymorphisms, rs1256049 and rs4986938. The expression of both these receptors has been recorded in testis and epididymis (8).

The aim of present study was to investigate association between ESR- α gene single nucleotide polymorphisms rs9340799 and rs2234693, and ESR- β gene single nucleotide polymorphisms rs1256049 and rs4986938 with implantation failure in Iranian northwest infertile women who were submitted for conventional in vitro fertilization (IVF) procedure and had no blastocyst implantation.

Materials and Methods

Collection of Patients and Samples

In this case-control study, we recruited 120 women (20-50 years old) referred to Medical

Genetics Laboratory, Tbriz, Iran, from July 2019 to Apr 2020. In this among, 60 IVF-treated women with implantation failure were recruited as case groups with confirmed idiopathic. Also, 60 IVF-treated women with successful implantation recruited as control group. The demographic information includes age, body mass index (BMI), alcohol drinking, tobacco smoking, and family history of case and control groups were collected by questionnaires and interviews from studied subjects (Table 1). All studied subjects were informed about the study and signed consent form, according to the Declaration of Helsinki ethical standards.

Primer Design

The ESR- α and ESR- β genes sequence were obtained from National Center for Biotechnology Information (NCBI) database. The Primer3 software were used to primer design for amplification of ESR- α gene rs2234693 and rs9340799 polymorphisms, and ESR- β gene rs4986938 and rs1256049 polymorphisms. The designed sequences of primers were synthesized in SinaClon Company, Iran. The sequences and characteristics of primers used amplify studied genes polymorphisms were presented in Table 2.

DNA Extraction and Genotyping

The peripheral blood samples (5 ml) were drawn from all subjects, and genomic DNA extraction was performed using salting out method. The quantity and quality of extracted DNA were investigated using nanodrop instrument and electrophoresis on 1% agarose gel. The genotype determination was performed by tetra-primer amplification refractory mutation system-polymerase chain reaction (Tetra-ARMS PCR) method. A 25 μ L total volume was used as following: PCR buffer (2.5 μ L), 1 μ L template DNA, 2.5 μ L dNTP (2 mM), 0.1 μ L each primers (25 pmol), 0.8 μ L Mgcl₂ (50 mM), and Taq DNA

polymerase (0.25 µL). PCR condition was as following: 1 cycle initial denaturation (94°C for 4 minutes), 40 cycles denaturation (94°C for 40 seconds), annealing (30 seconds), and extension (72°C for 25 seconds), and 1 cycle final extension (72°C for 5 minutes). The amplified PCR products were electrophoresed on 1% agarose gel with 50bp size marker. A gel documentation instrument was used to visualize the bands of PCR products.

Statistical Analysis

The statistical analysis was performed using SPSS software (version 19.0). The association of gene rs2234693, rs9340799, rs4986938, and rs1256049 polymorphisms

and implantation failure was evaluated by logistic regression. Moreover, the chi-square (χ^2) test and Fisher's exact test were used to evaluation of Hardy-Weinberg equilibrium (HWE). The difference between clinical and demographic features between healthy controls and patients were analyzed using independent sample t-test. The statistically significant was considered as $p < 0.05$.

Results

The obtained results showed no significant difference between patients and healthy controls in terms of family history, alcohol drinking, age, body mass index (BMI), and tobacco smoking (Table 1).

Table 1. The clinical features and demographic variables of cases and controls.

| Variables | Patients (n=60) | Controls (n=60) | P value |
|-------------------------------|-----------------|-----------------|---------|
| Age (year) | 34.19 ± 1.20 | 32.34 ± 3.76 | 0.659 |
| Body mass index (kg/m) | 25.09 ± 4.09 | 23.18 ± 3.11 | 0.398 |
| Tobacco smoking | | | 0.118 |
| Never | 49 (81.66%) | 52 (86.66%) | |
| Ever | 11 (18.33%) | 8 (13.33%) | |
| Alcohol drinking | | | 0.326 |
| Never | 41 (68.33%) | 43 (71.66%) | |
| Ever | 19 (31.66%) | 17 (28.33%) | |
| Family history | | | 0.211 |
| Negative | 56 (93.33%) | 58 (96.66%) | |
| Positive | 4 (6.66%) | 2 (3.33%) | |

Data are shown as mean ± SD or number (percent).

According to the χ^2 tests, the rs9340799 A/G, rs2234693 T>C, rs1256049 G>A, and rs4986938 G>A polymorphisms was in HWE in women with implantation failure and healthy controls ($p > 0.05$). The genotypes and alleles frequencies in women with implantation failure presented in Table 3.

We observed a significant difference between case and controls in term of rs9340799, rs2234693, rs1256049 polymorphisms; whereas rs4986938 polymorphism showed no significant association between case and healthy controls (Table 3).

Moreover, the frequency of C allele of rs2234693 in patients and healthy controls were 34% and 18%, respectively. Also, the frequency of A allele of rs1256049 in patients and healthy controls were 76% and 45%, respectively. The statistical analysis of C allele (rs2234693) and A allele (rs1256049) frequencies showed a significant difference between patients and healthy controls ($p > 0.05$); whereas alleles frequencies of rs9340799 and rs4986938 polymorphisms showed no significant association between case and healthy controls (Table 3).

Table 2. The sequences and characteristics of used primers.

| Gene | Polymorphism | Primer sequence | Genotype |
|---------------|--------------|---|---------------------|
| ESR- α | rs2234693 | Forward Outer: 5'-ACCACCATGCTCAGTCTCTAC-3' | 553+379 bp (TT) |
| | | Reverse Outer: 5'-CAAAACATGCACTCTCTGGGA-3' | 221+553 (CC) |
| | | Forward Inner: 5'-ATCTGAGTTCCAAATGTCCCATCT-3' | 221+379+553 bp (CT) |
| | | Reverse Inner: 5'-GAAACAGAGACAAAGCATAAAGCG-3' | |
| | rs9340799 | Forward Outer: 5'-ACCACCATGCTCAGTCTCTAC-3' | 262+553 bp (AA) |
| | | Reverse Outer: 5'-CAAAACATGCACTCTCTGGGA-3' | 330+553 (GG) |
| | | Forward Inner: 5'-CAGAGACCCTGAGTGTGGTATG-3' | 262+330+553 bp (AG) |
| | | Reverse Inner: 5'-CAATGCTCATCCCAACGCT-3' | |
| ESR- β | rs4986938 | Forward Outer: 5'-GTATGACCTGCTGCTGGAGA-3' | 191+486 bp (GG) |
| | | Reverse Outer: 5'-ATTGCGAGCCCTTCCAAGT-3' | 333+486 (AA) |
| | | Forward Inner: 5'-TGGCCACAGAGGTCAAAA-3' | 191+333+486 bp (GA) |
| | | Reverse Inner: 5'-CTGGAGTTCACGCTTCATCC-3' | |
| | rs1256049 | Forward Outer: 5'-GGCAGCCAAGCATCAACATTCTCAG-3' | 360+552 bp (GG) |
| | | Reverse Outer: 5'-ATTGCAGCACCCAGGACTTTGTTC-3' | 198+552 (AA) |
| | | Forward Inner: 5'-GGAGCTCAGCCTGTTTCGACCAATTA-3' | 198+360+552 bp (GA) |
| | | Reverse Inner: 5'-ATCCAACAGCTCTCCAAGAGCAGC-3' | |

ESR: estrogen receptor alpha; bp: base pair

Table 3. Genotype and allele distribution of ESR- α and ESR- β genes polymorphisms.

| Gene | Polymorphism | Genotype and Allele | Patients (n=60) | Controls (n=60) | OR (95% CI) | P value |
|---------------|-----------------|---------------------|-----------------|-----------------|---------------------|---------------|
| ESR- α | rs9340799 (A>G) | AA | 15 (25%) | 36 (60%) | Ref =1 | Ref |
| | | AG | 42 (70%) | 18 (30%) | 1.003 (0.108-1.645) | 0.001* |
| | | GG | 3 (5%) | 6 (10%) | 1.342 (0.229-1.664) | 0.14 |
| | | A | 72 (60%) | 90 (75%) | Ref =1 | Ref |
| | | G | 48 (40%) | 30 (25%) | 1.022 (0.963-2.284) | 0.54 |
| | rs2234693 (T>C) | TT | 21 (35%) | 39 (65%) | Ref =1 | Ref |
| | | TC | 37 (62%) | 21 (35%) | 1.233 (0.130-2.372) | 0.003* |
| | | CC | 2 (3%) | 0 (0%) | 1.332 (0.988-1.235) | 0.32 |
| | | T | 79 (66%) | 99 (82%) | Ref =1 | Ref |
| | | C | 41 (34%) | 21 (18%) | 1.544 (0.887-1.356) | 0.001* |
| ESR- β | rs1256049 (G>A) | GG | 3 (5%) | 30 (50%) | Ref =1 | Ref |
| | | GA | 23 (38%) | 6 (10%) | 1.943 (0.199-2.874) | 0.001* |
| | | AA | 34 (57%) | 24 (40%) | 1.123 (0.984-1.454) | 0.23 |
| | | G | 29 (24%) | 66 (55%) | Ref =1 | Ref |
| | | A | 91 (76%) | 54 (45%) | 1.107 (0.121-1.118) | 0.005* |
| | rs4986938 (G>A) | GG | 45 (75%) | 48 (80%) | Ref =1 | Ref |
| | | GA | 13 (22%) | 12 (20%) | 1.989 (0.112-2.322) | 0.344 |
| | | AA | 2 (3%) | 0 (0%) | 1.170 (0.567-1.777) | 0.19 |
| | | G | 103 (86%) | 108 (90%) | Ref =1 | Ref |
| | | A | 17 (14%) | 12 (10%) | 1.656 (0.340-1.544) | 0.32 |

*Statistically Significant; OR: Odds Ratio; CI: Confidence Interval

Discussion

Even in the best assisted reproductive technology units, pregnancy rate is not more than 30% (9). Different factors may be involved in failed pregnancy during the cycle, such as inappropriate ovarian stimulation, unsuitable uterus milieu and suboptimal laboratory culture conditions

(10). Implantation failure is the most common cause of lack of pregnancy after embryo transfer in IVF procedure. It takes place in about of 40% of IVF experiments (11). Some evidences indicate that genetic factors regulate implantation process. Thus, genetic defects and polymorphisms influence increasing susceptibility to process failure both directly and indirectly (12, 13). Among

the factors involved in the pregnancy, estrogen hormone has a crucial role. Fetal growth and development during both extra- and intra-uterine periods of life depends on estrogen produced by placenta (14). As it was mentioned before, null female mice with knockout ESR1 gene are infertile, with no corpus luteum formation and altered gonadotropin levels (15).

Considering the pivotal role of estrogen and its receptors during pregnancy, some studies have been done to find out whether there is any association between ESR- α and ESR- β variants and factors involved in pregnancy. The single nucleotide polymorphisms rs9340799, rs2234693, rs1256049, and rs4986938 is possible that it can influence ESR- α and ESR- β genes expression and subsequently pregnancy fate. In a study by Liaqat et al. reported that ESR α (rs2234693 and rs9340799) polymorphisms were strongly associated with risk of infertility in Pakistanian population; whereas ESR β (rs1256049) polymorphism no association with infertility risk (16). In another study by Ganesh et al. suggested that ESR α gene rs2234693 and rs9340799 polymorphisms are involved in increase of implantation failure in IVF-treated women in Indian population (17). Zulli et al. showed that ESR β gene rs4986938 polymorphism can be associated with risk of infertility in Brazilian women (18). Moreover, it has been revealed that ESR- α and ESR- β polymorphisms have association with ovarian response to follicle stimulating hormone in a person who undergoes IVF procedure (19). Based on an investigation in Turkish infertile women regarding IVF parameters such as numbers of collected oocyte, maturation and embryo quality, it has been suggested that these parameters could be associated with estrogen receptors variants (20).

In this study we tried to find an association between XbaI variant in ESR1 gene and blastocyst implantation failure in women

who undergo conventional IVF cycles. Our results showed that there is some association though not significant between the two. Effect of ESR1 gene variants over expression level of ESR1 gene may be useful to clear the potential role of these variations, but the exact mechanism behind the association remains unclear so far (21).

The reasons for the difference of reported results among the above mentioned studies could be due to several other related genes and environmental factors, and difference in studied samples size, ethnicity, race, and geographical area (22-25).

Considering the high medical cost and mental burden of couples who suffer from infertility, doing genetics screening along with clinical screening seems necessary before using artificial techniques. Beside other genomic markers, high valuable single nucleotide polymorphisms that have a significant association with infertility can be promising to avoid unpleasant results from failure cycles.

Conclusion

Generally, our study indicated more details of implantation failure, and suggested that rs9340799 and rs2234693 (ESR- α) and rs1256049 (ESR- β) polymorphisms may be associated with implantation failure in the Iranian northwest population. On the other hands, our study showed no association between rs4986938 (ESR- β) polymorphism and implantation failure in Iranian northwest population. Nevertheless, to reach the better understanding about implantation failure, in view of its multi-genetic background, identification of gene variants would determine the treatment strategies of the subjects.

Acknowledgments

This article was extracted from the MS.c. project of Parisa Pagard where Dr. Seyyed Ali Rahmani, advised and Lida Heidari advised

this project (Code: 10230503971013). This study was supported by Islamic Azad University, Tabriz, Iran. We thank the whole staff of Dr. Seyed Ali Rahmani medical genetics laboratory for assistance in the

successful strategy of this research.

Conflict of Interests

The authors declare no conflict of interest.

References

1. Paria BC, Reese J, Das SK, Dey SK. Deciphering the cross-talk of implantation: advances and challenges. *Science*. 2002;296(5576):2185-8. doi: 10.1126/science.1071601.
2. Shiralizadeh J, Barmaki H, Haiaty S, Faridvand Y, Mostafazadeh M, Mokarizadeh N, et al. The effects of high and low doses of folic acid on oxidation of protein levels during pregnancy: a randomized double-blind clinical trial. *Horm Mol Biol Clin Investig*. 2017;33(3):20170039. doi: 10.1515/hmbci-2017-0039.
3. Ma WG, Song H, Das SK, Paria BC, Dey SK. Estrogen is a critical determinant that specifies the duration of the window of uterine receptivity for implantation. *Proc Natl Acad Sci USA*. 2003;100(5):2963-8. doi: 10.1073/pnas.0530162100.
4. Ponglikitmongkol M, Green S, Chambon P. Genomic organization of the human oestrogen receptor gene. *EMBO J*. 1988;7(11):3385-8. doi: 10.1002/j.1460-2075.1988.tb03211.x.
5. Enmark E, Peltö-Huikko M, Grandien KA, Lagercrantz S, Lagercrantz J, Fried G, et al. Human estrogen receptor β -gene structure, chromosomal localization, and expression pattern. *J Clin Endocr Metab*. 1997;82(12):4258-65. doi: 10.1210/jcem.82.12.4470.
6. Ogawa S, Lubahn DB, Korach KS, Pfaff DW. Behavioral effects of estrogen receptor gene disruption in male mice. *Proc Natl Acad Sci USA*. 1997;94(4):1476-81. doi: 10.1073/pnas.94.4.1476.
7. Keene KL, Mychaleckyj JC, Smith SG, Leak TS, Perlegas PS, Langefeld CD, et al. Comprehensive evaluation of the estrogen receptor α gene reveals further evidence for association with type 2 diabetes enriched for nephropathy in an African American population. *Hum Genet*. 2008;123(4):333. doi: 10.1007/s00439-008-0482-z.
8. O'donnell L, Robertson KM, Jones ME, Simpson ER. Estrogen and spermatogenesis. *Endocr Rev*. 2001;22(3):289-318. doi: 10.1210/edrv.22.3.0431.
9. Allen VM, Wilson RD, Cheung A, Blight C, Désilets VA, Gagnon A, et al. Pregnancy outcomes after assisted reproductive technology. *J Obstet Gynaecol Can*. 2006;28(3):220-33. doi: 10.1016/S1701-2163(16)32112-0.
10. Isazadeh A, Azimian SH, Tariverdi N, Rahmani SA, Esmaeili M, Karimkhanlouei S, et al. Effects of coagulation factor XIII (Val34Leu) polymorphism on recurrent pregnancy loss in Iranian Azeri women. *J Lab Med*. 2017;41(2):89-92. doi: 10.1515/labmed-2017-0012.
11. Dentillo DB, Souza FR, Meola J, Vieira GS, Yazlle ME, Goulart LR, et al. No evidence of association of MUC-1 genetic polymorphism with embryo implantation failure. *Braz J Med Biol Res*. 2007;40(6):793-7. doi: 10.1590/S0100-879X2007000600007.
12. Nasirpour H, Key YA, Kazemipour N, Majidpour M, Mahdavi S, Hajazimian S, et al. Association of rubella, cytomegalovirus, and toxoplasma

- infections with recurrent miscarriages in bonab-Iran: A case-control study. *Gene Cell Tissue*. 2017;4(3):e60891. doi: 10.5812/gct.60891.
13. Isazadeh A, Hajazimian S, Rahmani SA, Mohammadoo-Khorasani M, Samanmanesh S, Karimkhanlouei S. The effects of Factor II (rs1799963) polymorphism on recurrent pregnancy loss in Iranian Azeri women. *Riv Ital Med Lab*. 2017;13(1):37-40. doi: 10.1007/s13631-017-0145-y.
14. Isazadeh A, Hajazimian S, Rahmani SA, Mohammadoo-Khorasani M, Moghtaran N, Maroufi NF. The effect of factor-xi (rs3756008) polymorphism on recurrent pregnancy loss in Iranian Azeri women. *Gene Cell Tissue*. 2017;4(1):e43717. doi: 10.17795/gct-43717.
15. Emmen JM, Korach KS. Estrogen receptor knockout mice: phenotypes in the female reproductive tract. *Gynecol Endocrinol*. 2003;17(2):169-76. doi: 10.1080/gye.17.2.169.176.
16. Liaqat S, Hasnain S, Muzammil S, Hayat S. Polymorphism analysis in estrogen receptors alpha and beta genes and their association with infertile population in Pakistan. *EXCLI J*. 2015;14:1085. doi: 10.17179/excli2015-559.
17. Ganesh V, Venkatesan V, Koshy T, Reddy SN, Muthumuthiah S, Paul SF. Association of estrogen, progesterone and follicle stimulating hormone receptor polymorphisms with in vitro fertilization outcomes. *Syst Biol Reprod Med*. 2018;64(4):260-5. doi: 10.1080/19396368.2018.1482030.
18. Zulli K, Bianco B, Mafra FA, Teles JS, Christofolini DM, Barbosa CP. Polymorphism of the estrogen receptor β gene is related to infertility and infertility-associated endometriosis. *Arq Bras Endocrinol Metabol*. 2010;54(6):567-71. doi: 10.1590/S0004-27302010000600010.
19. Boudjenah R, Molina-Gomes D, Torre A, Bergere M, Bailly M, Boitrelle F, et al. Genetic polymorphisms influence the ovarian response to rFSH stimulation in patients undergoing in vitro fertilization programs with ICSI. *PLoS One*. 2012;7(6):e38700. doi: 10.1371/journal.pone.0038700.
20. Ayvaz ÖÜ, Ekmekçi A, Baltacı V, Önen Hİ, Ünsal E. Evaluation of in vitro fertilization parameters and estrogen receptor alpha gene polymorphisms for women with unexplained infertility. *J Assist Reprod Genet*. 2009;26(9-10):503-10. doi: 10.1007/s10815-009-9354-2.
21. Bretherick KL, Hanna CW, Currie LM, Fluker MR, Hammond GL, Robinson WP. Estrogen receptor α gene polymorphisms are associated with idiopathic premature ovarian failure. *Fertil Steril*. 2008;89(2):318-24. doi: 10.1016/j.fertnstert.2007.03.008.
22. Soheilyfar S, Nikyar T, Fathi Maroufi N, Mohebi Chamkhorami F, Amini Z, Ahmadi M, et al. Association of IL-10, IL-18, and IL-33 genetic polymorphisms with recurrent pregnancy loss risk in Iranian women. *Gynecol Endocrinol*. 2019;35(4):342-5. doi: 10.1080/09513590.2018.1528220.
23. Hajizadeh YS, Emami E, Nottagh M, Amini Z, Maroufi NF, Azimian SH, et al. Effects of interleukin-1 receptor antagonist (IL-1Ra) gene 86 bp VNTR polymorphism on recurrent pregnancy loss: a case-control study. *Horm Mol Biol Clin Investig*. 2017;30(3):20170010. doi: 10.1515/hmbci-2017-0010.
24. Fathi Maroufi N, Gholampour Matin M, Ghanbari N, Khorrami A, Amini Z, Haj Azimian S, et al. Influence of single nucleotide polymorphism in IL-27 and IL-33 genes on breast cancer. *Br J Biomed Sci*. 2019;76(2):89-91. doi: 10.1080/09674845.2018.1545554.

25. Maroufi NF, Aghayi E, Garshasbi H, Matin MG, Bedoustani AB, Amoudizaj FF, et al. Association of rs1946518 C/A Polymorphism in Promoter Region of

Interleukin 18 Gene and Breast Cancer Risk in Iranian Women: A Case-control Study. Iran J Allergy Asthma Immunol. 2019;18(6):671-78.